

CORRECTION OF LEG DEFORMITIES
AND RESTORATION OF
FUNCTION OF LEG BONES BY

ILIZAROV TECHNIQUE

Edition - II



A TECHNICAL MANUAL FOR
ORTHOPAEDIC SURGEONS

MD.MOFAKHKHARUL BARI

ISBN: 978-0-9967956-1-6



This Technical Manual is written for professional orthopaedist and traumatologist mastering the method of transosseous osteosynthesis by Ilizarov technique.

Aim:

The aim of this manual is to highlight the main principles of treatment of patients with an orthopaedic pathology of leg bones (deformities, shortenings, defects, pseudoarthroses) using the method of transosseous compression-distraction osteosynthesis with the Ilizarov fixator.

Professor A.V. Gubin,

MD, MS, Ph.D., D.Sc.
Director, RISC, RTO, Kurgan – Russia

Academician Professor V.I. Shevtsov

MD, MS, Ph.D., DSc
Former Director, RISC, RTO, Kurgan – Russia,
Corresponding Member, Russian Academy of Medical Sciences.

Author:

Md. Mofakhkharul Bari
MD, MS (Kiev), Ph.D.(Tashkent)
Post Doctoral Fellow, Russian Ilizarov Scientific centre, Kurgan, Russia
Visiting and Honored Professor,
Russian Ilizarov Scientific Centre (RISC, RTO), Restorative Traumatology and Orthopaedics, Kurgan, Russia

Published by:

MedCrave Group LLC
2nd Edition – July 18, 2015

Foreword



It is indeed a great pleasure for me to write a few words on the eve of the publication of this excellent book titled “Correction of Leg Deformities and Restoration of Function of Leg Bones by Ilizarov Technique”, by our Honored Prof. of Kurgan Ilizarov Centre Md. Mofakhkharul Bari.

Bangladesh is a developing country and is burdened with a huge number of people with disabilities in desperate need of attention. But unfortunately many of the present day Orthopaedic surgeons are not well equipped with a knowledge to provide the treatment to these patients. Infact, many of the standard Orthopaedic books do not provide information for the young surgeons. So, a need for this kind of book has long been felt. Prof. MM Bari, a teacher with distinct quality and a lifelong devotee of deformity correction has given time and effort to bring out this book. This book is extremely informative and is most up-to-date. It is very stylishly written and neatly designed. What makes the book is that it makes the reading very interesting and thought provoking. The surgeon will find the book very gripping and absorbing.

My heartiest congratulations to him on his solo Herculean efforts. I wish him all the success.

A. V. Gubin
02.06.14

Prof. A. V. Gubin

MD; MS; Ph.D; D.Sc.
Director, Russian Ilizarov Scientific Centre, Restorative Traumatology and Orthopaedics (RISC, RTO), Kurgan, Russia.

Foreword



This book “Correction of leg deformities and restoration of Function of Leg Bones by Ilizarov Technique” is designed to provide the Orthopaedic surgeons with a rational treatment of trauma and orthopaedic problems. Professor Md. Mofakhkharul Bari’s large volume of personal expertise with Ilizarov technique since '82 (during Orthopaedic residency) till date is a wonderful achievement in the field of trauma and Orthopaedic surgery. This book characterizes the professional career of Professor M. M. Bari. I have watched Professor M. M. Bari silently grow over the 32 years to become a leading author in the field of Ilizarov surgery. This book is very informative and thought provoking. Simple writing, lucid language, clarity of thought, innovative technique, clinical photographs, good radiographs are all there in plenty. Prof. Bari was my very favourite fellow student. I congratulate Prof. M. M. Bari on his stupendous efforts and this is undoubtedly a great pride and honour for him. He has truly put Bangladesh on the world map and deserves praise and appreciation for all his efforts. I consider Professor Bari very successful. I hope and believe that this publication will be well accepted by the readers at home and abroad.

I wish him all the best and feel privileged to be able to write a foreword for this beautiful book.

V. I. Shevtsov 2.06.2014

Academician Prof. V. I. Shevtsov

MD; MS; Ph.D; D.Sc.
Former Director, Russian Ilizarov Scientific Centre, Restorative Traumatology and Orthopaedics (RISC, RTO), Kurgan, Russia
Corresponding Member
Russian Academy of Medical Sciences

Foreward to the Second Edition



I am honored and humbled to have this opportunity and share with you my deepest inspirations for writing a few words regarding the book titled "Correction of Leg deformities and restoration of function of leg bones by Ilizarov Technique". I must congratulate Professor M. M. Bari for his unique and specialized second edition who is the most experienced orthopaedic surgeon in the field of Ilizarov surgery. Ilizarov external fixation is ideal for open wounds, soft tissue infections, fractures and dislocations, osteomyelitis, deformities and poor circulations. In all these scenarios Ilizarov can be used through limited incisions and still providing stabilization, compression or off-loading the soft tissue envelope. Ilizarov compression distraction apparatus as it applies to the leg become widely accepted for trauma, reconstruction, lengthening and limb salvage procedures. The advantages conferred including use in the presence of infection and gradual ability to correct deformities appear to be a logical application of the technology. Large gaps any deformities, osteomyelitis with Ilizarov eliminates any windows and can be successfully applied during any condition of the limb. The use of gradual controlled post-operative correction with Ilizarov can achieve the same goal with minimal soft tissue handling. With the principles and techniques, the only limit for applications as applied to the leg is the surgeons imagination. With proper pre-operative algorithm the hope is that all surgeons can successfully apply an Ilizarov fixator consistently and in a timely manner. This book is comprehensive, simple and neatly written. A good book is one which apart from convincing interest in the students about the subject, makes them desirous to know more and more about it. This book does that and I am sure my Orthopaedic students will enjoy reading it. It is indeed be fitting that I write a forward for this book. Prof. M.M. Bari is a good trendsetter as well as orthopaedic writing is concerned and is worthy of emulation.

I wish him all the best.



Prof. (Dr.) Shamsuddin Ahmed

MBBS, FCPS, FRCS, FA Ortho, FICS
Formerly Professor of Orthopaedic Surgery
National Institute of Traumatology, Orthopaedic & Rehabilitation,
Dhaka Orthopaedic and Accident Surgeon First Recipient of
"Arthur Eyre-Brook Gold Medal" Awarded By World Orthopaedic
Concern

Message



It is a great privilege and honor for me to write a few words on the auspicious occasion of the publication of “Correction of Leg Deformities and Restoration of Function of Leg Bones by Ilizarov Technique” by Md. Mofakhkharul Bari who is a devotee to Ilizarov Surgery.

Our center, Bari-Ilizarov Scientific Centre (BISC) where he has done most of his magical works is a well recognized orthopaedic center making difference in the lives of human being. Ilizarov method has now gained a strong foothold in Bangladesh by which we can correct any congenital, acquired deformities and limb lengthening and solve any trauma problems. The motto of our center is to correct deformities, make people able to do their normal work and go back to their normal life.

My appreciation and congratulations to Dr. MM Bari for his tireless work in publishing this book.

A handwritten signature in blue ink, followed by the date 02/06/15.

Nabia Bari

MSS. IR (International Relations) Managing Director
Bari-Ilizarov Orthopaedic Center

Acknowledgements

It is a token of respect to my patients without whom I could not do this “**Correction of Leg deformities and restoration of function of leg bones by Ilizarov Technique**” which cannot be expressed in a few pages.

I have learned a great deal from my mentor prof. V. I. Shevtsov (former Director RISC, RTO, Kurgan, Russia) is my beloved teacher, Prof. A B Gubin, Director, RISC, RTO are the leaders of overall advancement in this field.

I recall the blessings given to me by Prof. E T Skliarenko (my MS supervisor, from Kiev) and Prof. B M Mirazimov (my Ph.D. supervisor, from Tshkent), because of them my career could get a new shape. My wife Nabia Bari could work on this modern gadgets and my loving sons and daughter and entire family have been the backbone in all my efforts, they really deserve appreciation. Finally I pray to Almighty Allah for the well-being of my patients.

Preface

I am greatly indebted to all my reverend teachers and fellow colleagues for helping me in writing this book while working in the then Soviet Union (Kiev Scientific Research Institute of Orthopaedics and Traumatology, ashkent Scientific Research Institute of Orthopaedics and Traumatology and Russian Ilizarov Scientific Centre) and different hospitals in Bangladesh from primary health care centres to the tertiary centres. I found that the fracture care has not gained much attention within the time of golden hour, that is the reason for development of deformity, non-unions and mal-unions which could have been prevented and corrected during the initial management at trauma hospitals. In our medical colleges, the orthopaedic units are not well equipped for managing the operation fractures and complex trauma patients, deformity, whatever maybe, is totally unproductive to the nation as a whole. The reconstruction of the limb deformity is my effort to bring attention towards it and fulfill the basic need of the hour in trauma care. Other techniques in orthopaedics, one can learn in 1-2 weeks; but one cannot learn this technique in a short period of time.

Many people have asked me “How did you get into Ilizarov technique?” I would like to acknowledge the mentorship and support of many people by telling the story of my orthopaedic Ilizarov journey. The most complex and post-traumatic problems were bone loss and large deformity and nobody could give a good solution for these. That’s why I have taken a decision to pursue fellowship in Kurgan (the mother institute) under the guidance of Prof. V. I. Shevtsov, the successor of Prof. G. A. Ilizarov and that was life changing for me. I was totally amazed by his clinical work. He is a great teacher and a great man and I am indebted to him for teaching me the tools of this amazing technique. In Kurgan I met N. M. Marzhuk, V. M. Shegirov, Y. P. Saldatov. I worked with them on clinical and academic pursuits and they exposed me to many aspects of the classic Ilizarov method.

I owe a great deal of gratitude to Prof. G. A Ilizarov, the great self-taught Orthopaedic surgeon, a pioneer genius with distinguished academic activities. This book “**Correction of Leg deformities and restoration of function of leg bones by Ilizarov ring fixator**” is based on the experience and treatment of patients for the last 32 years- in abroad, MMCH, Narayanganj 200 bedded hospital, Bio-Centre and NITOR. I believe that this manual will stimulate budding doctors to explore this sub-specialty orthopaedic surgery. I am the President of ASAMI, Bangladesh Chapter at present. I always encourage doctors who are interested in this field to become the member of this association. It is a place where we can learn, share ideas and meet fellow “Ilizarovians”. This manual comprises of clinical photographs accompanied by brief captions wherever it is useful, by operating pictures, diagrams, tables and charts. I offer my sincere thanks to all who have directly or indirectly helped me in the preparation of this manual in the last few years. I am thankful also to the staff in NITOR and BIO-Centre for their help at all times. My wife Nabia Bari (MSS, IR) who despite her family responsibilities and hospital Managing Director load has not only contributed but also helped in editing this manual, and our children Shayan Bari, Zayan Bari and Ishmam Bari have stood beside me. I am greatly indebted to my patients for providing me with both typical and atypical problems to study. The first step is the pre-operative planning which is universally required and beneficial. The typing, setting up manuscripts and photographs is the real tough job in preparing a manual and I appreciate the kind contribution by my eldest son Shayan Bari, Prof. Md. Shahidul Islam (Anaesthesiologist), Dr. Nazmul Huda Shetu, Dr. Mahfuzer Rahman, Hossain, Shahjahan, Munir, Jamir of Bio-Centre.

I wholeheartedly always welcome readers comments and criticism ideas and after all appreciation too. Thisbook is dedicated to my parents, my father and mother-in-laws and to my patients.


02/06/14

Md. Mofakhkharul Bari

MD, MS, PhD, Post-Doctoral Ilizarov fellow, Kurgan, Russia
Visiting and Honored Professor
Russian Ilizarov Scientific Centre, Kurgan, Russia. President ASAMI, Bangladesh

Preface to second edition:

The purpose of this book is to instruct the orthopaedic surgeon in the methods of applying the Ilizarov fixator. The book demonstrate how to apply the Ilizarov apparatus in common orthopaedic disorders. My purpose is to familiarize the orthopaedic and trauma surgeon who performs or intends to perform Ilizarov surgery with the myriad of parts, components and techniques required for successful treatment.

Professor Ilizarov was always fond of saying "No surgeon is justified to do any reconstructive surgery by Ilizarov technique unless he or she has got the competent idea regarding the technique and the apparatus". So, I feel one should know the basic principles of Ilizaov technique, the hardware, proper selection of patients, safe corridors of the placement of the Ilizarov's wire. Understanding of basic Kinesiology and Kinematics, anatomy and basic physiology are mandatory before attempting to master this Ilizarov technique. Development in Orthopaedic surgery in the past 3 decades have indeed remarkable particularly so with regard to compression-distraction method. Reconstruction by Ilizarov technique is an art and skill, which can only be mastered by the discipline of observation and practice. Ilizarov method has now gained a strong foot hold in Bangladesh.

The principles of Ilizarov surgery is based on thorough technical preparation that integrates both biological and biomechanical principles. Special attention to the patient's need in the post-operative period is absolutely mandatory to nurture the healing process.

The success of this method is now well established throughout the world for solving the most difficult problems of the musculoskeletal system. We have our own vast experience in this field which require a deep understanding of the method over a steep learning curve. This method cannot be learned from the lecture alone, and many details of each operative intervention and follow up must be critically reviewed.

In kurgan, RISC, RTO, the orthopaedic Mecca, I have witnessed a growing revolution in orthopaedics and traumatology, where and how biological approach for regenerating body parts has been born. Why replace body parts when we can regenerate them with living host tissue? Now time has come for whole world to return from metallurgy to biology. The biological insight's gleaned from Ilizarovs research have how been reproduced experimentally; but the learning curve is steep. Our recommendation is novice begin with simple lengthening, non unions and bone transports in the tibia only after doing a fellowship on Ilizarov technique. Later on you can proceede in the different part of the limbs.

This book was an instant hit among the students and orthopaedic surgeons justifying my efforts for the change. Every orthopaedic surgeon spoke highly about this book and this increased my responsibility for fulfilling the best expectations for the second edition. I have done a thorough overhauling of the book. While maintaining the basic things and have made the following changes in this edition.

- **More external fixation system have been added to beautify the book and make it more effective.
- **Color diagrams and new x-rays have been added to the text.
- **Each and every chapter has been updated and revised.
- **Orthopaedic Glossary has been added.
- **Pioneers in orthopaedics and their contributions have been duly acknowledged.

With all these features I have made a very sincere effort to cover up the deficiencies. I am very much grateful to all my students and teachers who took the trouble of writing to use their opinions and I have tried to incorporate all the healthy suggestions given by them. I believe the Second Edition will be equally appreciated by all concerned.

I wholeheartedly welcome readers' comments and criticisms. This book is dedicated to my dear parents, father and mothers-in-law and to all my patients.

I appreciate the kind contribution of my sons Shayan Bari & Zayan Bari & my daughter Ishmam Bari; and Md. Masudur Rahman of IT Section, Bari-Ilizarov Orthopaedic Centre.

For me it has been both an honor and privilege to work with my beloved mentor Academician Professor V.I. Shevtsov in orthopaedics in editing the book. I was amazed by Professor V.I. Shevtsov's clinical work. The Kurgan fellowship with Professor V.I. Shevtsov and others was life changing. They are great teachers and I am indebted to them for teaching me the tools of this amazing trade. I sincerely hope that the work presented here will receive the attention of trauma surgeons as well as orthopaedic surgeons. I share the philosophy that Ilizarov technique is still the best treatment option for all kinds of deformities to restore bone and limb alignment as well as to reconstruct joint congruence not only in young but also in specific groups of older patients. With limb preserving treatments limb anatomy can be reconstructed preventing secondary osteoarthritis of joints.

২০/০৩/২০১৫

Md. Mofakhkharul Bari

MD, MS, PhD, Post-Doctoral Ilizarov fellow,
Kurgan, Russia

Visiting and Honored Professor

Russian Ilizarov Scientific Centre, Kurgan, Russia. President ASAMI, Bangladesh

Former Unit Chief, Limb Lengthening,

Surgical Reconstruction and Deformity Correction Unit National Institute of Traumatology and Orthopaedic Rehabilitation (NITOR)

Sher-E-Bangla Nagar, Dhaka, Bangladesh.

Contents	
Chapter I	1
History.	
The Ilizarov Method: History	
Chapter II	5
The Michelangelo of Orthopaedics.	
Chapter III	
Memorable Moments	7
Honored Professor Ceremony in Ilizarov Centre, Kurgan, Russia, 14 June, 2013	
Chapter IV	12
Development of Compression-Distracton Method and External Transosseous.	
Fixation of Bones	
Chapter V	18
Basic Ilizarov Frame and its Components.	
Chapter VI	28
Anatomo-Physiological Peculiarities of Children Extremities in using	
Compression-Distracton Method.	
Chapter VII	30
Classification of Compression-Distracton Method.	
Chapter VIII	32
General Principles of Application of Apparatus.	
Chapter IX	35
Ilizarov Basic Concepts:	
Proper Selection of Entry and Exit Points for Passing Ilizarov's or K-Wires	
Proximal tibial wires	
Middle tibial wires	
Distal tibial wires	
Hybrid technique	
Middle tibial Schanz Screw	
Distal (Lower) tibial Schanz Screw	
Calcaneal Wires	
Lateral Approach	
Proper position of the rings	
Half Calcaneal ring	
Forefoot half ring	
Equinus and Equinovarus with short limb (Equinus with LLD)	
Osteotomy of fibula	
Corticotomy of tibia	
Anatomical factors	
Biomechanical factors	
Physiological factors	
Common mistakes during corticotomy	
Limb lengthening	
Biology	
Distracton Histogenesis of Soft Tissues	
Distracton is indicated for	

Complications of limb lengthening	
Joint complications	
Soft tissue complications	
Post operative correction and management	
Chapter X	44
Hormonal Changes in Limb Lengthening Process.	
Chapter XI	46
Blood Supply and Metabolic Activities of Bony Tissue.	
Chapter XII	48
Biochemical Procedure of Metabolism in Lower Limb Lengthening in Adult Group.	
Enhancement of Regenerate Bone Healing.	
Distraction osteogenesis can be characterized into three groups	
Motorized distraction	
HBO (Hyperbaric Oxygen Therapy)	
Anticatabolic therapy	
Chapter XIII	50
Mechanical Principle of Ilizarov Method.	
Chapter XIV	54
Different Plane (Frontal plane, Sagittal plane and Horizontal plane) Deformities and Biomechanics of the Ankle and Foot.	
Frontal / coronal plane deformities of ankle and foot	
Sagittal plane deformities of ankle and foot	
Arches of the foot	
Different planes and deformities	
Chapter XV	58
Disorders of the Bone Configuration and Function of Leg Bones.	
I. Deformities of Leg Bones	
II & III. Pseudoarthroses and Defects of Leg bones	
IV & V. Leg Lengthening, Leg Shortening with Segmental Axis Disturbance Leg Shortening without its Axis Disturbance	
Reconstruction of the Limb Versus Amputation.	
Case Study 1	
Case Study 2	
Case Study 3	
Open Fracture Tibia and Fibula.	
Case Study 4	
Mal Uniting Fracture Right Lower Tibia Fibula (3 & ½ months old) GII.	
Case Study 5	
Open Fracture Right Lower Tibia Fibula G-III A.	
Case Study 6	
Infected non Union Right Tibia Fibula with Interlocking Nail in situ GIIIB, Nail is Exposed.	
Case Study 7	
Infected Open Fracture Right Tibia Fibula GIIIB with Exposed Plate & Screws.	
Case Study 8	
Open Fracture (GIIIA) Left Tibia Fibula.	

Case Study 9
Open Comminuted Fracture Left Lower Tibia Fibula (GII).
Case Study 10
Comminuted Fracture Right Lower Tibia Fibula.
Case Study 11
Open Segmental Comminuted Fracture Left Lower Tibia GII.
Case Study 12
Non Union Left Upper Tibia.
Case Study 13
Aseptic Nonunion of Long Bones with Anterior Bowing.
Case Study 14
**Infected Gap non Union Right Lower Tibia Fibula, Equinus Foot with AO Fixator in situ,
Treated in Apollo Hospital, Dhaka.**
Case Study 15
Gap Nonunion of Left Tibia with Gross Deformity and LLD (12.6).
Case Study 16
Non Union Left Femur Due to Gunshot Injury, 13 cm LLD.
Case Study 17
Infected Gap non Union Right Lower Tibia, Deformed Foot with Biaxial Fixator *in situ*.
Case Study 18
Osteomyelitis of Right Upper Tibia Leading to Gap non Union.
Case Study 19
Chronic Osteomyelitis of Left Tibia.
Case Study 20
Non Union of Right Lower Tibia with Broken Nail.
Case Study 21
Post Traumatic Left Tibia Vara.
Case Study 22
Post Traumatic Tibia Vara with Bad Scar.
Case Study 23
Right Anteromedial Bowing.
Case Study 24
Tibialization.
Case Study 25
Case Study 26
Post Osteomyelitic Big Gap non Union of Right Tibia.
Case Study 27
Hockey Stick Deformity (C-Type Deformity).
Case Study 28
Anterolateral Bowing.
Case Study 29
Left Anteromedial Bowing.
Case Study 30
Rt Anteromedial Bowing.
Case Study 31
Antero Lateral Bowing of Left Tibia.
Case Study 32

Congenital Shortening of Left Tibia.

Case Study 33

Congenital Shortening of Left Tibia (4.8 cm LLD) with Absence of 2 Digits and Short Foot.

Case Study 34

Normal Height Increase of an Individual Using Ilizarov Technique.

Case Study 35

Normal Height Increase.

Case Study 36

Normal Height Increase.

Case Study 37

Post- Polio Residual Deformity - 5.3 cm Shortening.

Case Study 38

Post Polio Residual Deformity - LLD 8 cm.

Case Study 39

Post Polio Residual Deformity - LLD 5 cm with Unstable Knee, Ankle Facing Forward, Equinovarus Deformity.

Case Study 40

Post Polio Residual Deformity, 4 cm LLD With Right Ankle Valgus.

Case Study 41

Post Polio Residual Deformity of Right Leg, Ankle and Foot with External Tibial; 2 cm LLD with Equinovarus >65o (Severe).

Case Study 42

Neurofibromatosis.

Case Study 43 & 44

Fibrous Dysplasia Right Tibia.

Case Study 45

Hypertrophic Nonunion of Tibia, Monkey Mouth Deformity.

Case Study 46

Post Traumatic Disorganized Knee, Ankle Valgus and 14 cm LLD.

Case Study 47

Mal Union Right Upper Tibia, Varus Deformity with 4 cm Shortening with External Rotation Accompanied by Equinus Deformity >45°.

Case Study 48

Mal Union Right Middle Tibia with Bowing Deformity.

Case Study 49

Genu Valgum Deformity.

Case Study 50 & 51

Chapter XVI

125

Author with Foreign Dignitaries.

Author as a Faculty in Different Countries of the Globe.

References

128

Glossary

129



Chapter I

History

History

Professor Gabriil Abramovitch Ilizarov– graduated from medical school in the Soviet Union in 1943 near the end of World War II. After graduation, he was assigned to practice in Kurgan, a small town in western Siberia. He was the only physician with hundreds of miles and had little in the ways of supplies and medicine. Faced with numerous cases of bone deformities and trauma victims due to the war, Professor Ilizarov used the equipment at hand to treat his patients. Through trial and error with handmade equipment, this self-taught orthopaedic surgeon created the magical combination that would cause the bones to grow again. Thus the Ilizarov technique was created.



Academician G.A. Ilizarov
15.06.1921 – 24.07.1992

Ilizarov Technique- The Ilizarov technique/method of treatment used Prof. Ilizarov's Principle of Distraction Osteogenesis. This refers to the formation of new bone between two bone surfaces that are pulled apart in a controlled and gradual manner. The distraction initially gives rise to Neo Vascularisation, which is what actually stimulates the new bone formation. In addition, there is histogenesis of muscles, nerves and skin and in diseases (osteomyelitis, fibrous dysplasia, pseudoarthroses) this new bone replaces pathological bone with normal bone. This is a revolutionary concept in medical science; diseases for which earlier there was no treatment possible can now successfully be treated with Ilizarov Method.



bone healing and immobilization of fractures. While he was studying Mechanics he had an insight into the stability that an external ring with crossed wires would bring to a fracture setting. He then asked a local metal worker to fashion these specially designed parts for a new orthopaedic device, and as before did his preliminary testing on a broken broom stick. This time Ilizarov became convinced that his invention would provide secure immobilization. He sent an application for a certificate of invention and was invited to Moscow to demonstrate his external fixator. This application for authorship of the device



Typical of Siberia the institute is shaped like a snowflake. Each wing houses a separate clinical service, sharing central facilities of operating room, Radiology, Laboratory and Physical Therapy.

was accepted in June 1952 and was finally approved in 1954. When he presented his data at the conferences, other physicians were skeptical because his results of treatment were dramatically shorter. That being said, devices similar to the Ilizarov apparatus started to emerge, in spite of the skepticism, using bow circular and rectangular shaped fixators. One of them was half ring external fixator by Gudushauri in 1954 from CITO. That was the "official" external fixator used in Moscow for many years. At that time Moscow "coryphaei" did not recognize "non-yielding" province director from Siberia. Prof Volkov was because the director of CITO in 1961 was one of the prominent figures who actively



Entry to Ilizarov's new hospital called the Russian Ilizarov Scientific Center, Restorative Traumatology and Orthopaedics (RISC, RTO inaugurated in 1983).

The Ilizarov Method: History

In 1950, Ilizarov moved to Kurgan. In Kurgan, Ilizarov continued to explore ways to achieve improved results in



worked against official acceptance of the Ilizarov device and method. In 1968 Prof. Volkov and Prof. Oganesyanyan had patented a similar device to the one presented by Ilizarov at the conference on TB of bones and joints in Tomsk in 1963. But Volkov used his prestige and position; promote the application of his device in the Soviet Union and at international conferences. However the fortunes of Dr. Volkov would dramatically shift during perestroika, when he was removed in 1985 from his position as director of CITO. Among the reasons for his dismissal was oppression towards the acceptance and distribution of Prof. Ilizarov's external fixator and his method.

Author in the dog square. Dogs are helping in research work.

In 1965, he was awarded the title of "Honoured physician of Russia Federation" for his achievements in medicine. He also became known among patients, title as the "Magician from Kurgan". The 'title' and little



Front Gate of the Ilizarov center.

In front of Ilizarov's monument author with Prof. A.V. Gubin, Director of the Ilizarov Institute.

to do with Ilizarov's lifelong love of hearing and showing off magic tricks, which became his hobby throughout his life and provided him great joy and relaxation [2-4].

Despite the negative experiences in Moscow and his first attempts to introduce his new device to the Soviet Union. Prof. Ilizarov and his innovative treatment started to gain recognition locally. In 1955 he became the chief of the department of Trauma and Orthopaedics in the Veterans hospital in Kurgan.

In 1965, the Health Ministry decided to send a group of physicians to Kurgan to observe more clearly the surgery and progress of the patients according to Ilizarov method. In 1968 Ilizarov operated on Valery Brumel (1942-2003), a famous Russian athlete renowned in international sports, as high jumper who had set six world records during the 1960s. Tragically, he suffered an open fracture of the distal tibia in a motorcycle accident. The accident and his injuries and treatment received wide notice in the Soviet press. Brumel spent 3 years in various clinics and underwent about 20 unsuccessful operations.

Ultimately, he developed an infected non-union as well as significant LLD. In 1965, Dr. Golyakhovsky [5] was among the First group of specialists sent to Kurgan to observe and evaluate Prof. Ilizarov's work. Dr. Vladimir [5]. Golyakhovsky [5] was a young successful surgeon from CITO (Central

Institute of Traumatology and Orthopaedics, Moscow). Dr. V. Golyakhovsky spent one month in Kurgan and had returned to CITO, Moscow. He was amazed about the treatment and methods of Ilizarov; but Prof. Volkov of CITO was apathetic about supporting the device within CITO.



In front of Ilizarov's museum author with Prof. A.V. Gubin, Director of the Ilizarov Institute. That chaika car was used by Ilizarov.

Meanwhile Valery Brumel was started to lose hope of recovery in CITO. Dr. Golyakhovsky advised him to go to Kurgan to seek a consultation with Prof. Ilizarov. On being reassured that his infected non-union could be healed as his LLD of 3.5 cm could be corrected. The surgery was successful. Brumel resumed his athletic training session in 1968. These events brought substantial recognition and attention of high officials as well as fame to Prof. Gabriel Abramovich Ilizarov within the Soviet Union. Brumel's recovery was also published in the U.S. Medical Press, the Journal of Podiatry (Foot disorders) in 1973 and was titled "Kurgan: Revolution in Orthopaedics".

These media exposure of a famed athlete recovering against such overwhelming odds and prior treatments failures helped to gather support for financing a new orthopaedic Institute in Kurgan in 1971 (KNIIEKOT).


In 1982 an additional building in the shape of a snowflake was added to expand the clinic, research and diagnostic services of the institute. This snowflake design was an original concept of Prof. Ilizarov to prevent the spread of infection, by placing patient wards furthest away from the administrative center of the building and providing them with independent entrances. In case of serious infection breaking out in one ward, that block could be effectively isolated without interruption of the work of the hospital.



G. A. Ilizarov with Valery Brumel during his treatment in Kurgan.



Author in the graveyard of Academician G.A. Ilizarov.

A large, light blue, stylized letter 'M' logo is centered on the page. The 'M' is composed of thick outlines and is superimposed over a faint, light blue globe. The globe shows continents and is partially obscured by the 'M'.

Chapter II


The Michelangelo of Orthopaedics

The Michelangelo of Orthopaedics

Carlo Mauri (1930-1982) who was a well known Italian journalist, alpinist and explorer, helped to introduce the Ilizarov system into Western Europe. In 1980 Professor Ilizarov did the surgery in the Mauri's infected tibial non union and which was completely healed. On his return to Italy Mauri wrote an article in an Italian newspaper, naming the Russian Physician Gavriil Ilizarov as the "Michelangelo of Orthopaedics". This would prove to be the break in the dam that would release the news of the Ilizarov method and external fixator to the world medical community and set in motion events that have led to world wide application and study of the Ilizarov method. Carlo Mauri's physician in Italy was amazed by the healing that had occurred of his longstanding non union condition. Mauri subsequently invited Prof. Ilizarov to Italy in June 1981 and arranged for his participation in XXII Italian AO conference in Bellagio, Italy through his physician friends. Italian physicians immediately realized the significance of these "Siberian technique" and were enthusiastic to learn the procedure.

American surgeons first learn this technique from their European colleagues. Dr. Frankel with Dr. Stuart Green visited the Ilizarov center in Kurgan in 1987. In 1987 Ilizarov visited New York and Dr. Golyakhovsky recall this event "Ilizarov requested three carousel projectors for his lecture, which surprised the inviting party. The auditorium was packed. People were sitting and standing in the aisles. Ilizarov showed 700 slides in one and half hours. When he finished the audience jumped from their seats and applauded, standing, for about 10 minutes. Interest in the Ilizarov external fixator in USA was contagious and many started to use the apparatus without proper training, making mistakes that led to complications as well as discouragement and misplaced blame on the external fixator. Ilizarov said "Boldness should not exceed one's skill". Fortunately, many of the orthopaedic surgeons carefully studied this method and became the world known famous expert.





Chapter III

Memorable Moments

Memorable Moments



Author in the Chamber of Academician G. A. Ilizarov Museum.



Author in the O.T of Academician G.A. Ilizarov Museum.



In front of Ilizarov center Kurgan.



Dog Square



Author is doing operation in Ilizarov center (August 2012) Prof G. P. Ivanov is assisting him.



Author with Honored prof. of Kurgan Ilizarov center S. I. Shevd.



Flags of different countries. Author in Ilizarov conference June 13-15 2013; Kurgan Russia.



Author with Prof. A. V. Gubin, Director (RISC, RTO) in front of Ilizarov monument.



Author with N. Murzhikov.



Author with Valentina Kamysheva (Inter. Head) and N. Murzhikov



Author with Prof. Y. P. Saldatov.



Dog Square, Dogs are helping in Scientific research



Author in the department N-10 during his visiting Prof. program July 2012



Professor V. I. Shevtsov (Former Director RISC, RTO) with Author's family (Ishmam Bari, Nabia Bari), Greece, 2012



Delegates from different countries; in front of Ilizarov center, Kurgan June 2013



Professor A. V. Gubin with Professor MM Bari and Shayan Bari Directors office, Kurgan, Russia.

Honored Professor Ceremony in Ilizarov Centre, Kurgan, Russia, 14 June, 2013



Welcome address by Prof. A.V. Gubin



Opening up the gown



Opening up the gown



Wearing the gown



Opening up the gown



Opening up the gown



Opening up the gown



The Moment of wearing the Cap



Cap on the head



Deputy Director for Scientific works Y. Borzhunov Deputy Director for Scientific works Y. Borzhunov is clipping broach of Honored Prof. to MM Bari. Prof. A. V. Gubin is also welcoming him.



Delegates are welcoming Prof. M.M. Bari for with huge applause being the Honored Prof. of RISC, RTO



Clipping the Honored Prof. Broach.



Deputy Director for Scientific works Borzhunov is congratulating Honored Prof. M.M. Bari



Prof. A. V. Gubin is welcoming Honored Prof. of Ilizarov center M. M. Bari




Author is presenting his book to Professor Y. P. Saldatov after the Honored Professor Ceremony



Honored Prof. of Ilizarov Center Prof. S.I. Shevd and Prof. M.M. Bari Cap



Y. Borzhunov Deputy Director for Scientific works, Professor A. V. Gubin with Honored Professor MM Bari and Honored Professor S I Shevd 14-06-2013



Professor A. V. Gubin, Honored Professor MM Bari, Professor V. I. Shevtsov (14-06-2013, 6:10 PM)

Chapter IV

Development of Compression-Distraction Method and External Transosseous Fixation of Bones

Development of Compression-Distraction Method and External Transosseous Fixation of Bones

Compression-distraction is an independent sector in orthopaedics and traumatology has got definite history of its own development and may be analyzed on many parameters. Utilization of this method in orthopaedics attracts many specialists with opportunity without considerable surgical trauma to eliminate severe congenital & acquired deformities, which is connected with great regeneration.

Undoubtedly facts remains that compression-distraction method opens new era instead of using traditional methods of surgical interventions. Compression-distraction and reposition-distraction apparatuses were developed by K.M. Sibash (1952); G.A. Ilizarov (1952); O.N. Gudushauri (1954); S.S. Tkachenko, V.K. Kalnberz (1971) and others. In 1952. G.A. Ilizarov developed a principle of new type of cross wires transosseous apparatus and it is consists of cross wires and metallic rings. Subsequently other apparatuses were developed by other scientists. Among these wide spreading were found the apparatus of G.A. Ilizarov; V.K. Kalnberz; O.N. Gudushauri; Volkov-Oganesyan. Methods of limb lengthening suggested by G.A. Ilizarov were universally recognized and widely used in practice. Good results using the transosseous osteosynthesis of Ilizarov for treating pseudoarthrosis, defects and deformity of bones confirms highly its effectiveness. Compression distraction method in most of the cases gives good results in treating pseudoarthrosis and defects with infection (Gudushauri O.N. 1964; Shumilkina E.I.; Matusis J.E. 1970; Tashpulatov A.G. 1985 and others). Development and improvement of this prespective method of treatment have several directions. Firstly, it creates more perfect opportunity and constructions. Secondly it brings different changes in construction and details that have in transosseous apparatus. Thirdly new method of treatment are developing in orthopaedotraumatological patients with the use of compression distraction method. In 1843 Malgaigne designed a special device for external fixation of bone fragments in patella and olecranon fractures (Figure 3.1). It consists of two plates, each of them ending with two hooks. A spacing screw connects the plates. When a pair of hooks is introduced through the skin into each fragment, the plates and hooks fixed in the bone fragments are drawn together until tight contact is made and therefore reciprocal fixation of the fragments.



Figure-3.1: Apparatus of Malgaigne.

In 1902 Lambotte designed an apparatus (Figure 3.2) to fix fractures outside the fracture site. The apparatus consists of screws tightly fixed in bone fragments (2 to 3 screws in each) and two plates connecting the protruding ends of the screws by means of bolts.




Figure-3.2: Apparatus of Lambotte.

In 1919 there were many publications on transfixion of bone fragments. Transfixion (Figure 3.3) means- a nail is introduced transversely into each fragment to fracture, distally and proximally, after reposition, then a circular plaster dressing is applied, the nail ends being cast into it.



Figure-3.3: Transfixion of bone fragments.

In 1925 the device designed by Rosen L., consists of a T-shaped plate with two slots at right angles to each other and two screws with two fixing nuts in each. After the screws are introduced into bone fragments, the T-shaped plate is put on the protruding ends of the screws (Figure-3.4). By shifting both screws along the slots and nuts along the screws, a surgeon shifts the fragments with respect to each other, thereby achieving reposition of the fragments.

In 1929 the device designed by Hempell consists of two semi rings, which fix the bone fragments with pins. The rings are connected to each other by screw rods.




Figure-3.4: Apparatus of Rosen.




Figure-3.5 : Apparatus of Hempell.

In 1934 Bittner developed his apparatus (Figure 3.6) which consists of pins, metal rings, and hinged distractors. After a pin is passed through each fragment, the ends were fixed in the rings. The rings with fragments fixed there in were drawn apart or together by means of the distractors connecting them, which resulted in distraction or compression of bone fragments.




Figure-3.6: Apparatus of Bittner.

In 1937 Stader designed an apparatus (Figure 3.7) that he applied successfully in dogs with various fractures. His device consist of a pair of pins for each fragment and an adjusting connecting bar. The pins are to be inserted into the fragments at an angle to each other. The fragments are drawn apart or together with the aid of a turnbuckle. A screw arrangement at each end of the connector helps to reduce angular displacement of bone fragments. Later on Stader "reduction and fixation splint" were regularly used in man.




Figure-3.7: Stader Splint.

In 1938 Petrovsky designed an apparatus (Figure 3.8) for bone lengthening. After osteotomy, a pin was introduced into each fragment, the pin ends were fixed in metal rings. Two turnbuckles attached to the connector bars served the purpose of drawing the rings end therefore the fragments, apart or together. The idea of Petrovsky apparatus was used in some other external transosseous fixators that were designed later.




Figure-3.8: Apparatus of Petrovsky.

In 1942, Hoffmann applied an apparatus (Figure 3.9) of his own design in various bone injuries. Several tip threaded pins are screwed into each fragment, either along a straight line or in an arbitrary pattern. The protruding tips of pin groups are fixed with clamps. Two threaded rods are hinged to the clamps and connected with a turnbuckle. After repositioning, the connecting rods are locked in their hinges, thus fixing the fracture. According to Hoffmann the apparatus often reduced fragments and fixed them firmly in a closed manner. To increase stability of the apparatus, Vidal added another frame to the apparatus.




Figure-3.9: Apparatus of Hoffmann with a Vidal double frame.

In 1944 Boston Pediatric Orthopaedic Surgeon Bost F. used the external transosseous fixation apparatus of his own design for the lengthening of lower extremity bones (Figure-3.10). For lengthening of femur, Z-osteotomy was followed by the insertion of two cross pins into the femur inferior to the lesser trochanter. The ends of those pins were fixed in the ring. Two other pins were passed through the lower third of the femur and fixed the ends of the pins in the lateral planks of the device. The ring and lateral planks were connected to each other with two sliding rods. The ring and planks with fixed bone fragments were disjoined using the nuts on the rods thus performing the bone fragments distraction. The rate of bone lengthening made up 3 to 4 mm daily. The apparatus was dissembled in 12-14 weeks and plaster cast was applied. That plaster cast was used to fix the ends of one upper pin and second upper pin that were kept in the extremity. Weight bearing was allowed 6 months after the operation. Plaster cast was used for 1 year. Application of that technique gave from 5 to 7 cm lengthening.




Figure-3.10: Apparatus of Bost.

In 1952 Sivash designed a fixator (Figure 3.11) that consists of two split screws with special clamps for pins, the basic principle is that the resected bones are rapidly drawn together with a force sufficient for their impacting. This accelerates bone union by several times.




Figure-3.11: Apparatus of Sivash.

In 1952 an apparatus (Figure-3.12) designed by Professor G. A. Ilizarov of Kurgan has come into wide use in Soviet Union. It is intended for arthrodesis, in diaphyseal fractures, non unions, pseudoarthrosis, limb lengthening and for deformity corrections. This device consists of rings and cross wires, the rings are connected by spacing screws. Two Ilizarovs wire are passed through each fragment crosswise, clamped to the metal ring and tensioned with a special appliance called Tensioner. 4 spacing screws provide for compression or distraction of bone fragments.




Figure-3.12: Apparatus of Ilizarov.

In 1953 Wittmosser R. invented an apparatus of his own design for bone fragments fixation. It consists of crossing pins and rings connected to each other with extensible distractors. It has a special screw device for repositioning bone fragments in width (Figure-3.13).




Figure-3.13: Apparatus of Wittmosser.

In 1954 Gudushauri designed an apparatus (Figure-3.14) for reposition and fixation of bone fragments and for bone lengthening. The apparatus consists of two pairs of coupled bows, a repositioning bow, and two spacing screws. One pair of the coupled bows holds two pins transfixing the distal fragment; the other holds the pins transfixing the proximal one. A repositioning bow is located between the coupled bows, providing for repositioning of bone fragments in width. The spacing screws correct displacement of bone fragments in length and provide their compression or distraction. The Gudushauri apparatus was widely used for compression osteosynthesis in non union and pseudarthrosis and for fragment repositioning with subsequent compression.




Figure-3.14: Apparatus of gudushauri.

In 1962 Grishin designed an apparatus (Figure- 3.15) for arthrodesis of the ankle joint. It consists of two uniform halves. Each half has a head with two 'legs' hinged to it. The legs are provided with screws and turnbuckles. When the ankle joint is exposed and the cartilage covering the tibia and the talus is removed, three metal pins 3.5mm in diameter are passed in the transverse direction; one through the distal tibia, another through the calcaneus, the third through the talus. Then the foot is put at a functional position and the device is applied, with the pins clamped on both sides. According to Grishin, the foot can be reliably fixed in the apparatus at any angle of flexion or extension with respect to the leg. To correct undesirable flexion, the anterior part of spacing screws are drawn together, as are posterior, to avoid extension. Varus or Valgus deformity of the posterior part of the foot is eliminated by exerting more effort in drawing together the lateral or medial pair of spacing screws, respectively. The disadvantage of Grishin device is that thick pins are to be applied and they may injure both soft tissues and bone.




Figure-3.15: Apparatus of grishin.

In early 1970 Kalnberz of Riga, Alma- Ata designed an apparatus (Figure-3.16) which consists of plastic rings connected by coil springs. The coils serve as threads for nuts, with clamps allowing for holding and adjusting the rings. For compression, the springs are extended and they provide for constant traction. When apparatus is applied for distraction, the springs are to be compressed. The coil springs can also bend, which facilitates correction of limb deformities.




Figure-3.16: Apparatus of kalnberz.

In 1971 Valkov- Oganesyanyan reposition hinged distraction apparatuses (Figure- 3.17) are widely applied for reposition and fixation of bone fragments in fractures and pseudarthrosis, for limb lengthening, for fragment fixation after various correcting osteotomies, for compression arthrodesis.



Figure-3.17: Apparatus of volkov-oganesyan.

In 1988 the CITO (Central Institute of Traumatology and Orthopaedics, MOSCOW), orthopedic surgeons described the rod compression-distraction devices MKT (Fig.-3.19) for the fixation of bone fragments in fractures of long tubular bones and fractures of the pelvic bones, as well as for the treatment of congenital deformities. The apparatuses provide easy access to the fracture wounds and early mobilization of patients.




Figure-3.19: KT Apparatus



Fig.-3.18: Pichkhadze.

In 1974 Pichkhadze R. suggested the device for transosseous osteosynthesis in intra-and periarticular fractures of long tubular bones in shoulder, elbow and knee joints (Fig.-3.18). The apparatus is intended for and fixation of fragments in the required position with simultaneous development of movements in the injured and adjacent joints.



Chapter V

Basic Ilizarov Frame and its Components

Basic Ilizarov Frame and its Components

Osteosynthesis with Ilizarov frame is achieved by securing the bone fragment to the external fixator with wires. Ilizarov surgeon can assemble the individual components in to any configuration which is needed for particular problem.

External transosseous fixation system offers many advantages over the internal fixation devices:

1. Compression can be maintained during the entire treatment period.
2. Fixation can be obtained without inserting hardware at the site of pathology; this is very much important in case of infected non-union or pseudoarthrosis.
3. Less traumatic than the implantation of an internal fixation device.
4. The compression-distraction device can be removed without an additional operation.

Components:

Main support components are:

1. Half rings (Figure 4.4)
2. Arches (Figure 4.7, 4.8)
3. Long Connecting plates (Figure 4.15)

Auxillary support components are:

1. Short straight connecting plates (Figure 4.15)
2. Curved plates (Figure 4.16)
3. Twisted plates (Figure 4.17)
4. Posts, Male and female (Figure 4.43, 4.44)
5. Wires and olive wires (Figure 4.34, 4.35, 4.36)
6. Buckles (Figure 4.37)
7. Threaded rods (Figure 4.12)
8. Telescopic rods (Figure 4.13)
9. Wire fixation bolt (Figure 4.21)

Additional connecting elements are:

1. Bushing (Figure 4.19)
2. Threaded sockets (Figure 4.18)
3. Washers (Figure 4.23)
4. Bolts (Figure 4.20)
5. Nuts (Figure 4.21, 4.22)
6. Hinges (Figure 4.39, 4.40)
7. Multiple pin Fixation clamp (Figure 4.45)

Half Ring:

It has got the mechanical resistance greater than 90 Kg/ Square mm and it is made up of titanium metal.

Sizes : 12 Viz 80, 100, 110, 120, 130, 140, 150, 160, 180, 200, 220, 240 mm inner diameter.

Hole diameter: 08 mm.

Space between holes: 04 mm.

Its ends are bent into ledges. This allows attachment of one half ring to another in the same plane.



Figure 4.1: Steel ring.



Figure 4.2: Carbon ring (Non radio lucent and radio lucent).



Figure 4.3: Connecting ring.



Figure 4.4: Carbon half ring.

5/8th Ring:

Sizes: 130, 150, 160mm inner diameter.

Advantages:

1. Facilitates joint movements
2. Wound dressing can be done easily
3. Myocutaneous flaps and large deep incision as in compartment syndrome it is useful



Figure 4.5: 5/8th ring.

Application of location: Upper third of thigh & upper third of humerus.



Figure 4.8: Radiolucent Italian Arch (Carbon Composite).

Disadvantages:

1. Do not bear load of tensioned wires
2. It can be used only with full rings.

FOOT RING:

These rings are used in hind foot and fore foot to give better space for Ilizarov or K-wires.




Figure 4.6: Foot ring.



Figure 4.9 : Russian Arch

T handle - Wrench for half pins or Schanz. Useful instrument to introduce half pins or Schanz into the bone with free hand.



Figure 4.10 : T handle.



Figure 4.11 : Special (Arch) for children; authors modification of M. Catagni arch.

Arch

Original Russian Arch

Sizes: 80-260 mm inner diameter and is used in the upper 1/3 rd of the thigh to secure wires placed through the proximal femur.



Figure 4.7: Italian Arch (Non radiolucent).

Italian Arch:

Sizes: 90 and 120 small and large.

Advantages: Slots and holes to secure tapered Schanz or half pins in multiple planes.

Threaded Rods:

Diameter: 6mm

Thread pitch: 1mm

Sizes: 40, 60, 80,100, 120, 150, 200, 250, 300, 350, 400 mm. It is used to interconnect rings and arches.



Figure 4.12 : Threaded rods.

Telescopic Rod

Advantages:

1. It increases frame rigidity when connecting rings or arches are greater than 150mm apart.
2. It has a threaded stud at one end and a perpendicular locking bolt at the other end to hold the threaded rod.

Sizes available: 100, 150, 250, 350, 400mm. It is a long hollow tube with its inner diameter larger than the outer diameter of any threaded rod.



Figure 4.13 : Telescopic rods.

GRADUATED TELESCOPIC ROD

Advantage:

It is used in lengthening and provides direct measurement.

Sizes available: 60, 100, 150, 200, 250 mm.

Figure 4.14:
Graduated
Telescopic rods.

Connection Plate

Variants:

1. Curved
2. Twisted
3. Straight
 - a) Short connecting plate
 - b) Long connecting plate
 - c) Connecting plate with threaded ends.

Thickness : 5 mm

Wide : 4 mm

Hole diameter : 7 mm

Twisted Plate

This plates are used to connect between the holes of vertical and horizontal planes.

Length : 45 mm, 65 mm 86 mm

2 holes 3 holes 4 holes



Figure 4.16 : Twisted Plate.




Figure 4.15 : Connecting Plates.

Curved Plate

These are used as extension to half ring arch for accommodating Ilizarov or K-wires.



Figure 4.17 : Curved Plate.

Threaded Socket

Length: 20, 40 mm External diameter: 10 mm

Both ends are threaded to accommodate bolts or connecting rods. Two perpendicular threaded holes are provided at the center on either side to connect bolts or rods as extension of threaded rods.



Figure 4.18 : Threaded Sockets.

Bushing

Size: 12mm, 24mm with one and two perpendicular holes. It is one mm wider than threaded rod. It moves over the threaded rod. It can be used as a spacer.




Figure 4.19: Bushing.

Connecting Bolt

Sizes available: 10,16,20, 30 mm. Thickness: 4mm.

They can bridge the distance from the rings to pins or wires conveniently.



Figure 4.20 : Connecting Bolt.

Nuts

Thickness: 3,5,6mm

3mm is used for locking nuts on hinges.

5mm is used for stabilizing all forms of frame construction. 6mm is used for connecting rods where compression/distraction is required.



Figure 4.21 : Nuts in slotted and cannulated bolts.

Wahser:

It is used to fill the space. 4 types are available.

1. Spacing washer.
2. Split locking washer.
3. Flat sided washer.
4. Slotted washer.



Figure 4.22 :Nut.



Figure 4.23 : Washer.

Wire Fixation Bolts:

It is used to secure the wires to support components at the holes. Types

1. Slotted.
2. Cannulated.

These are of 6mm threaded diameter, 18mm length and a bolt head thickness of 6mm. The bolt heads are either hexagonal or with two flat and rounded surfaces. The cannulated bolts have a 2mm hole while slotted bolts have an oblique groove on the under surface of the bolt head.



Figure 4.24 : Wire fixation bolt slotted.

Wire Fixation Cannulated Bolt:

Cannulated bolts are preferred for 1.5mm wires and slotted bolts for 1.8 mm wires.



Figure 4.25 : Cannulated wire fixation bolt.

Block for Half Pins / Schanz

Sizes: 1,2,3,4 holes.



Figure 4.26: Block for half pins/Schanz.

Box Wrench for Bolt:

Is used to tight & loosen the nut.



Figure 4.27 : Box wrench for bolt.



Figure 4.28 : Spanner 10mm size.

Oblique Support:

is used to connect the Italian arch with ring.

Direct Measuring Wire Tensioner



Figure 4.29 : Oblique support



Figure 4.32 : Direct Measuring wire tensioner (Russian Dynamometer)




Figure 4.33 : Tensioning using wire tensioner.



Figure 4.30 : Different Box Wrenches.

Corticotome:

Sizes available: 3, 5, 7, 9 mm.



Figure 4.34 : Different corticotome.

Original Ilizarov Mechanical Wire Tensioner:



Figure 4.31 : Wire Tensioner Mechanical

Wires:

They serve to connect bones or bone fragments to the support elements of frame. They differ in diameter, length and shape of the point.

Bayonet point:

Advantages:

1. It has a greater penetrating power.
2. It causes less heating effect of bones and soft tissues.
3. It produces hole of a diameter slightly larger than that of wire, causing less friction.

Bayonet point cortical 1.5mmx300mm for forearms & foot bones.

Bayonet point cortical 1.8mm x 370mm for diaphysis of tibia and femur.



Figure 4.35 : Bayonet point.

Trocar point:

Advantages:

1. It is used for metaepiphyseal region because it has less penetrating power.
2. It has greater hold in the bones.
3. Trocar point cancellous 1.5 mm x 300 mm for metaepiphyseal region of radius & ulna
4. Trocar point cancellous 1.8 mm x 370 mm for metaepiphyseal region of tibia and femur.



Figure 4.36 : Trocar point.

Olive or Bead or Stopper Wire:

The olive or kink in the wire is used to achieve and maintain the position of fragment after the fracture. It has also a great role for deformity correction. Wires with olive are used in cases where much force (upto 180-200 kp) is applied to the bone. In osteoporotic bone; it is desirable to use wires with flat surfaced cones or wires with a corkscrew bend. The wire must be resilient with a well polished surface.




Figure 4.37 : Olive or bead or stopper wire.

Wire Fixation Clamp:

Buckle:

The buckle has a U-shaped configuration with short threaded rods on each arm of a short plate. The clamp is completed by employing a short two hole plate and a pair of nuts, which creates a rectangle. This clamp has a groove for a wire and a threaded hole to secure components. A

wire put into the groove of the clamp can be fixed to a support component - either a ring or a plate by tightening a small two hole plate against the supports opposite surface. The buckle is most commonly used where it becomes necessary to fix additional wires to The configuration.



Figure 4.38 : Buckle.



Figure 4.39 : Use of buckles to fix additional wires.



Figure 4.40: Hinges.



Figure 4.41: Formation of hinges.



Figure 4.42 : Female and male post.



Figure 4.43 : Universal joint and male post. hinges. Indications.

1. To correct any type of deformity.
2. It is used as a pivot (rotation) point component which is essential for strengthening.

Advantages:

1. Gives constrain motion in a specific plane.
2. It provides specific fulcrum for control of specific correction of angulations.

Formation of Hinges:

1. One male half hinge and one female half hinge.
2. Two female half hinges are connected to threaded rods.

Important parameters for positioning hinges:

1. Two rings to which hinges are attached must be strictly perpendicular to the bone fragments.
2. Two hinges are located at opposite sides of the deformity for stabilization.

3. The hinge rotation axis must be situated at the apex of horizontal level of the deformity. If two hinges applied both should be on the same level.
4. Hinges are applied at the same level of deformity.
5. The position of hinges can be used to achieve different types of deformity correction such as opening wedge distraction, compression, translation and derotation.

Speed of correction with hinges:

Rule of triangle: Speed of distraction, compression is transferred to the hinge axis by factor 3:1. i.e. for 1 mm movement at axis there must be 3mm movement at device site.

Post:

Posts are of 2 types. Male posts are 28,38 or 48 mm long with 2,3 and 4 holes respectively.

Female posts are 30, 40 or 50 mm long with 2, 3 and 4 holes




Figure 4.46: Multiple pin fixation clamp.



Figure 4.44: Male Post.



Figure-4.47: Schanz conical (different sizes).



Figure 4.45: Female Post.



Figure-4.48: Schanz tapered (different sizes).

Modification of oblique support by the author

Goniometer



Figure-4.49: Wire cutter.



Figure-4.50: Author's modified oblique support.



Figure-4.52: Different types.



Figure-4.53: Ilizarov box for different components.



Figure-4.51: Half rings; Variants of assembly.

Chapter VI

Anatomo-Physiological Peculiarities of Children Extremities in using Compression- Distraction Method

Anatomo-Physiological Peculiarities of Children Extremities in using Compression-Distracton Method

Anatomical and plastic characteristics of osseo-articular systems for children significantly differs from the adult. Most of the methods for treating the children is based on using these peculiarities and that is why is not used in adults. Child is born without ossified skeleton. Bones of the new born extremities consists of bony and cartilagenous parts. In children the periosteum is thick and loosely attached to the cortex and produces new bone rapidly. In adult periosteum is thin, adherent to the cortex and produces new bone less rapidly. For these fundamental differences, healing of fractures in children is rapid. Epiphyseal plate cartilage is responsible for the growth of bone in length. It receives nutrition from bone through epiphyseal and metaphyseal vessels.

Epiphyseal plate cartilage has got 4 layers:

1. Zone of resting cartilage.
2. Zone of proliferating cartilage.
3. Zone of maturing cartilage.
4. Zone of calcifying cartilage.

Ossification of cartilagenous parts of bones goes on gradually and with the growth of children it is finished at the age of 16-17 years. In its development osseoarticular system passes through 5 stages (Sadofev-V.I.1990).

1st stage (from birth to 6-10 months): Epiphysis and greater part of metaphysis of long bones have got cartilagenous structures. Diaphysis consists of little bony tissue.


2nd stage (from 6-10 months to 4 years): determines as a beginning stage of epiphyseal ossification of long bones. In this stage complete ossification of metaphysis occurs and part of the epiphysis also. Apophysis remains cartilagenous.

3rd stage Formation of osseo-articular system (from 4 years to 8-9 years): In this period complete ossification of epiphysis occurs. Cartilagenous structures at the end of 3 stages remains in the acromial end of clavicle, apophysis of all bones and growing zones.

4th stage: post natal formation of osseo- articular systems- epiphyseal and tuberosities of long bones and foot ossification occurs. Ages are from 15 to 18 years.

5th stage: post natal formation of osseo- articular systems-synostosis of metaphyseal and apophyseal growing zone occurs. Tentative ages are from 15 to 18 years.

Knowledge of development of stages and formation of osseo-articular systems in children have got great practical significance in finding indications towards different methods of compression-distracton treatment. In selecting the methods of operations it takes into account the anatomical peculiarities of bone for every concrete ages. For example wide use of distracton epiphyseolysis method demands the presence of well ossification and solid epiphysis, consequently in 1st and 2nd stage of development, it cannot be used. Compression-distracton method is also ineffective in the 5th stage of development of osseo-articular system due to formation of synostosis of growing zones. Proceed from anatomical characteristics, every stages of bone formation may conclude that compression- distracton method can be used from the middle of 2nd stage (1.5-2 years), when diaphysis and metaphysis of long bones ossifies and have definite stability.



Chapter VII

Classification of Compression - Distraction Method

Classification of Compression - Distraction Method


All methods for the treatment of orthopaedic pathology with compression-distraction method conditionally can be divided into 2 groups.


1. Apparatus methods of treatment: In this group those methods are included, that are not needed extrasurgical intervention in bones and soft tissue of extremities after applying the Ilizarov's apparatus. We can include here methods of correction of soft tissue deformity, contracture (CTEV, joint contracture etc) and distraction epi and metaphysiolysis.
2. Apparatus and surgical methods of treatment: This group includes those methods, when with the application of Ilizarov as well as surgical intervention is done in soft tissues and bones. Moreover, these interventions can be done in the same sitting with the application of Ilizarov (Osteotomy) for achieving the good outcomes (e.g. arthrodesis, skin grafting).

Orthopaedic intervention in the extremity can be classified in relation to the above mentioned classification

- a) Lengthening, shortening and widening of bones.
- b) Correction of bowing deformities
- c) Transportation and filling the defects of bones.
- d) Correction of contracture, dislocations and arthrodesis of joint.

Stable transosseous osteosynthesis is the basic optimum process for adaptation of organism in trauma as well as in surgery. There is a dilemma regarding the question of how much limb lengthening is possible in one stage surgery, however; literature informs lengthening can be achieved upto 24cm in one stage surgery.





Chapter VIII

General Principles of Application of Apparatus

General Principles of Application of Apparatus

Assembly:

The Assembly and design of the frame depends on the pathology, anatomic location and other features of local tissues. Rings provide rigid bone fragment fixation. Arches are used in locations where rings are either impossible to use or would interfere with joint motion: the hip, the shoulder, & the elbow.

Precautions:

Ring or arch should be about 1.5 to 2cm (at least 1 finger breadth) larger than the maximum diameter of the limb at each level of fixation.

General rules of Ilizarov application:

Biomechanical principle of Ilizarov ring fixator is very important for osteoneogenesis. That is why we must follow the some fundamental rules for better results:

1. The bone fragments must have the same position with respect to their rings or arch supports.
2. Connecting threaded rods must be parallel to each other and to the longitudinal axis of the bone fragments.
3. The wires must be tensioned uniformly, it should be maintained till consolidation of bone fragment is complete
4. Stable fixation is absolutely mandatory for pain free motion.
5. Use smaller rings for better stability and use larger rings whenever necessary.
6. Wire inserted close to the joints should not limit the range of motion.

Ilizarov wires insertion technique:

For each individual case we emphasize the following things into consideration:

1. The number of wires used.
2. Their positions and locations.
3. And the planes in which they are inserted.

The rigidity of fixation of bone fragment depends on both biomechanical and biological factors.

The mechanical factors are:

- a) The number of wires.
- b) The stiffness of wires.
- c) The shapes, size, and position of the rings.
- d) The distance between the wires.
- e) Other supporting elements of the frame.

The biological factors are:

- a) The level of fracture (Osteotomy)

- b) The position of the fracture plane.
- c) The stiffness of interfragmentary soft tissues.
- d) The width and contact surface area of the fragment ends.
- e) Local biologic status.

Wires can be inserted at any level of long bones. The direction and crossing of the wires is determined by the local anatomic constraints - the vessels, nerves and tendons. Wire should be inserted straight through the soft tissues down to the bone. Drilling should be started after the wire tip has engaged the bone surface.

Precautions:

To prevent thermal injury to the bone and soft tissues we must follow the following rules during drilling:

1. Cool the wire by holding it with an alcohol - soaked gauze or betadine gauze or hexisol gauze at the wire skin interface.
2. Give pause during drilling to allow the wire tip to cool.
3. Drive the tip of the wire quickly through the soft tissues on the same side of the limb.
4. Cool the wire tip with an alcohol gauze or betadine gauze, or hexisol gauze as it emerges.
5. When the bone is subcutaneous, like in medial surface of tibia insert, wires from the opposite side to prevent heat conduct direct to the skin.
6. Don't bend the wire while drilling; because deformed wire increases the size of the wire tract, traumatizes soft tissues, deviates from its path, can enlarge bone holes which reduces the rigidity of fixation and increases the risk of wire tract infection.
7. Transmedullary wire insertion is important rather than only cortical, because it provides wire stability and prevents cortical osteomyelitis.
8. The Close the wire is to the joint, more attention should be paid.
9. Introduction of wire must be done correctly at first attempt, the rule is, one wire one hole.

Soft tissue considerations:

To reduce skin necrosis at wire entry and exit sites during ring movement i.e., compression- distraction, angulations or transport by creating an excess stock of soft tissues at the appropriate location. If a compression osteosynthesis is planned, the skin and soft tissues are pulled away from the center of the configuration when inserting wires.

When distraction osteosynthesis or lengthening is done the stock of soft tissues is shifted slightly towards the center of the configuration at the time of wire insertion. Proper wire placement can preserve joint motion. When you insert a wire near a joint, extend the joint as a wire penetrates soft tissues on the flexor surface, and flex the joint as a wire passes through the extensor muscle.

Alignment of rings and arches:

As a rule, the position of the fragments with respect to their respective rings must be similar. If we can follow this principle, displaced fracture fragments will automatically reduce as the rings are made coaxial to each other. If the rings are made parallel the angular deformities will be eliminated.

Tensioning and securing wires:

After fixing the ring or arch in proper relationship to the bone's longitudinal axis, the assistant should hold the ring in position and the surgeon applies tension by tensioner and secures the wires. Each wire must remain straight as it is fastened to the ring.

Precautions:


Secure one end of the wire and apply tension to the other and try to maintain equal tension on the wires. This will create an even distribution of the forces on the apparatus. We must emphasize special attention to wires attached to arches. All the wires connected to an arch must be equally tensioned.


Maintaining wire tension:

We should maintain uniform wire tension during whole period of treatment. Sometimes we can use one or two supplementary wires through each bone fragment to improve stability in cases of compression osteosynthesis if the bone fragment ends are either incongruent or have large gaps along the edges.

Arrangement of connecting rods:

Stable fixation is absolutely mandatory for good regeneration. The rods should be parallel to each other and to the longitudinal axis of the bone segments which is especially very important for a frame that is used for compression, distraction or bone transport.





Chapter IX

Ilizarov Basic Concepts

Ilizarov Basic Concepts

In this Atlas our basic aim is to describe the use of Ilizarov technique for correcting the equinus and equinovarus and CTEV deformities of foot and ankle. For excellent outcome 3 important things have to be kept in mind.

1. Meticulous follow up.
2. Appropriate selection of rings and wires and other components.
3. Proper construction of frames prior to application.

Proper Selection of Entry and Exit Points for Passing Ilizarov's or K-Wires.

1. Initially if lengthening of tibia is needed along with deformity correction then in this situation three sets of cross wires are passed in the leg- the proximal, middle and distal.
2. Secondly, the frame configuration is indicated where only the correction of deformity is required. In this situation two sets of wires are passed at about the junction of proximal to the middle third and middle to the distal third.

Proximal tibial wires:

Entry and exit points

The first wire (1.8mm) enters through the most prominent part of the head of the fibula with an inclination aiming parallel to knee joint line and to emerge at a point just posterior to the vertical line drawn from the medial border of the patella. The second wire enters on the anterolateral surface of upper end of tibia just behind and about 1/2 cm (5mm) below the point drawn by the horizontal line of the head of the fibula and vertical line drawn from the lateral border of the patella with an inclination of the wire aiming to emerge at the same level just anterior to the medial border (Figure 5.1). Sometimes a 3rd wire can be passed from the lateral side anterior to the head of the fibula aiming to emerge on the medial side at the same level, in case of more stability and for tibial lengthening.



Figure-5.1: Proximal tibial wires.

Middle tibial wires:

Entry - Exit points

The first wire is inserted at about 1 finger breadth (2 cm) lateral and behind the shin of the tibia with an inclination of the wire aiming to emerge at the same level just anterior to the medial border. The 2nd wire is inserted




Figure-5.2: Middle tibial wires.

about 1 finger breadth (2 cm) anterior to the palpable anterior border of the fibula at a point about 1/2 cm (5 mm) below the level of the first wire with an inclination anteriorly aiming the wire to emerge about one finger breadth (2cm) postero- medial to the shin of tibia (Figure-5.2).

Distal tibial wires :

Two Ilizarov's wires or K-wires (1.8 mm) are inserted at the junction of the distal and middle third of tibia. The first wire inserted through fibula and passed through the tibia with an inclination aiming the wire to emerge one finger breadth (2cm) postero medial to the shin of the tibia (Figure-5.3). Regarding the insertion of the 2nd wire, one must be very much cautious. The space between the anterior border of the tibia and fibula is divided into three equal zones. * The anterior zone contains neurovascular structures and tendons. * The middle zone is safe and contains muscular parts. Now the 2nd wire inserted through safe corridor of middle zone 1/2cm (5mm) below the 1st wire into the tibia with an inclination to emerge just anterior to the medial border of the tibia (Figure-5.4). One must be careful to avoid piercing the Vena saphena magna (Great saphenous vein) which crosses the medial border of the tibia about 10cm (5 finger breadth) above the medial malleolus (Figure-5.5, 5.6, 5.7).



Figure-5.3: Entry point of distal tibial wires.




Figure-5.4: Exit point of 1st and 2nd wires.




Figure-5.7 : Position of wires (Middle and distal tibial rings).



Figure-5.5: Middle tibial wires.




Figure-5.8a: Proximal schanz screws.




Figure-5.6: Position of wires tibial rings (Proximal, middle, distal).




Figure-5.8b: Proximal schanz screws.




Figure-5.9a: Middle schanz screws.




Figure-5.9b: Schanz in lower 1/3rd of tibia

Hybrid technique:

Means the product of a cross between genetically unlike individuals i.e the combination of transosseous wires with Schanz or half pins.

The half pins inserted on the sagittal plane crosses the transverse wire at 90, allows for stability without transfixion of muscular masses. Half pin fixation or Schanz has the advantages of inserting at the site away from the neurovascular structure.

Proximal Tibial Schanz Screw: After inserting the 1st reference wire through the most prominent part of the head of the fibula with an inclination to emerge at a point just posterior to the vertical line dropping from the medial border of patella, one Schanz screw is passed through the anterolateral surface of upper end of tibia from a point about 1.2 cm. below the reference wire on the vertical line drawn from the lateral border of the patella. The screw is inserted with a inclination to engage the thicker cortex just behind the medial border (Figure-5.8a & 5.8b).

Middle tibial Schanz Screw:

One Ilizarov wire or K-wire and one Schanz screw is passed (Figure-5.9a).

Distal (Lower) tibial Schanz Screw:

The first Ilizarov or K-wire is inserted through the fibula and passed through tibia with an inclination aiming the wire to emerge one finger breadth (2 cm) postero- medial to the shin of tibia. First Schanz screw is fixed from medial border of tibia and the second Schanz screw from medial surface of tibia making maximum angle at a distance from 1st screw with the help of 3 to 4 hole Ranche Cube block (Figure-5.9b).

Proper points of Selection of wires in foot:

1. For calcaneum - 1.8mm two wires.
2. For mid-trasal - 1.5 mm two wires.
3. For metatarsals - 1.5 mm two wires.

Calcaneal Wires:

Medial approach:

The pulsation of posterior tibial artery should be felt and posterior tibial nerve can also be palpated. One and a half finger breadth behind artery and nerve is the safe corridor on the medial surface of calcaneum which emerges about 2 to 3 finger breadth (4-6 cm) behind the posterior border of posterior malleolus. The first K-wire is inserted at one finger breadth (2 cm) above the undersurface of calcaneum and one finger breadth (2 cm) in front of Tendo Achilles insertion. The second wire is inserted at a point 1cm anterior to the first point and 1cm above the inferior surface of calcaneum. The angles between the two wires will be more or less 30 degree (Figure- 5.10, 5.11, 5.12)



Figure-5.10: Wire position in calcaneus and metatarsals.




Figure-5.11: Wire in calcaneum.



Figure-5.12: Wire position in calcaneus and metatarsals.

Lateral Approach:

Insertion Point:

First wire is inserted through outer surface of calcaneum 1 cm above its inferior surface and 1 cm distal to the attachment of Achille's tendon and the wire is further introduced with an inclination anteromedially aiming to emerge on the medial surface of the calcaneum 2 finger breadth (4 cm) below the medial malleolus. The second wire is inserted one finger breadth (2 cm) distal to the entry point of first wire and is advanced posteromedially to emerge one finger breadth (2 cm) behind the first wire in the same level. Here the angle between the two wires should be more or less 30 degree (Figure-5.11 & 5.12).

Proper position of wires in mid tarsal region:

If equinus is associated with cavus two additional K-wires or Ilizarov's wire are required to correct such deformity. One wire is passed on the medial surface of navicular to emerge at the summit of the cavus and other wire is introduced at the centre of outer surface of cuboid and aiming to emerge in the summit of the cavus and both the wires are making angle of 30 degree between them. Wires should not be tensioned.

The first wire is inserted from the outer side of distal part of 5th metatarsal between the head & neck and is advanced obliquely piercing the 5th, 4th & 3rd metatarsal just proximal to their head to emerge on the surface on the dorsum of the foot. The second wire is inserted on the medial surface in the distal first metatarsal just proximal to its head and advances obliquely to emerge on the dorsum of the foot after piercing through second metatarsal.



Figure-5.13: Position of wires in mid tarsal region.

Position of wires at the base of metatarsal head:

The first wire is inserted from the outer side of distal part of 5th metatarsal between the head & neck and is advanced obliquely piercing the 5th, 4th & 3rd metatarsal just proximal to their head to emerge on the surface on the dorsum of the foot. The second wire is inserted on the medial surface in the distal first metatarsal just proximal to its head and advances obliquely to emerge on the dorsum of the foot after piercing through second metatarsal.




Figure-5.14a: Position of wires in mid tarsal region.



Figure-5.14b: (Position of K wires in standard approach).



Figure-5.14c: (Position of K wires in standard approach).
Proper size of the ring:

Try to use smaller rings but use larger rings whenever necessary. At least two finger breadth (3 to 4cm) should be considered at the maximum girth of the limb. Special attention should be given for proper selection of optimum measurement of the ring. But often the doctor selects the size of the ring by mere assumption his eyes. On the basis of our long years of clinical experience we can use transosseous apparatus through a mathematical calculation and we have identified the optimum measurement of rings, which depends on circumference of the extremity. For longer diameter rings it is better to use Ilizarov's wire Optimum measurement of ring always depends upon the volume of the extremity.

Circumference of extremity in cm	Measurement of Ring
10	90-100
15	100-110
20	120-130
25	140-150
30	150-160
35	160-180
40	180-195

Proper position of the rings:

Proximal tibial ring:

Proximal tibial ring is used when lengthening is performed. A full ring should be fixed at the level of the head of the fibula (Figure-5.15a & 15b).

Middle tibial ring:

A full ring is placed approximately at the junction of upper 1/3rd to middle 1/3rd of the leg (Figure-5.15C).

Distal tibial ring:

A full ring is placed about 10-12 cm above the ankle joint, about the junction of lower 1/3rd and middle 1/3rd of leg.




Figure-5.15c

Half Calcaneal ring:

Important thing is that the half calcaneal ring should be placed behind and parallel to plantar surface of heel, which will be more or less horizontal when the patient stands but in cases of varus or valgus deformity the inclination of the ring will be according to the concerned deformity and will not be horizontal (Figure-5.16).



Figure-5.16: Placement of the ring in calcaneum.

Forefoot half ring :

Here half ring is placed proximal to the head of metatarsals and the ring should be perpendicular to the head of metatarsals.




Figure-5.17: Placement of rings in calcaneus & forefoot.

Full ring in forefoot:

It is justify to mention that if the equinus deformity is combined with the cavus or excavatus a full ring should be placed around the forefoot (Figure-5.18).




Figure-5.15a & 5.15b: Position of rings in tibial lengthening.




Figure-5.18: Placement of half ring in calcaneus and full ring at forefoot.

Half ring in mid-foot:

A half ring is also sometimes needed in the mid foot when equinus is associated with cavus or excavatus deformity.



Figure-5.19: Placement of half ring in mid foot region.

Full ring in mid foot:

In the mid foot a full ring is also needed when equinus is associated with severe equinocavus deformity (Figure-5.20).




Figure-5.20: Placement of full ring in mid foot and forefoot region.

Pre construction of the frame:

To save time during surgery we can pre-construct the tibial frame. The frame constructed will be different for correcting equinus deformity alone and equinus deformity along with lengthening of short leg (Figure-5.21). For correcting

equinus deformity alone proximal ring is not at all required, middle & the distal ring will serve our purpose.




Figure-5.21a,b: Position of the rings for tibial lengthening.




Figure-5.22: Complete frame assembly two anterior and two posterior threaded connecting rods.

Equinus and Equinocavus with short limb (Equinus with LLD):

Equinus with associated short limb can be corrected simultaneously by lengthening the tibia in the metaphyseal region with fibular osteotomy.



Figure-5.23: Corticotomy of tibia

Osteotomy of fibula:

Osteotomy of fibula is performed at the junction of middle third and lower third of fibula. Osteotomy of fibula can be performed in two ways.

1. By making 2 or 3 drill hole over the fibula.
2. Or by direct using 1-1.5cm corticotome over the fibula.

Corticotomy of tibia:

Step 1

The levels of osteotomy, K-wires, knee joint space, fibula osteotomy should be determined and clearly marked.

- P - Patella.
- T.T - Tibial tuberosity.
- C.L - Corticotomy level.
- J.L - Knee joint line.

Step 2

Skin incision 2cm on the level of tibia osteotomy (0.5-1cm below tibial tuberosity).

- * Ilizarov corticotomy is a low energy osteotomy with the preservation of periosteum and endosteum.
- * In choosing corticotomy level, anatomic, biomechanical and physiological factors must be considered.

Anatomical factors:

Corticotomy should not be performed in the middle of the bone.

Corticotomy is suitable in between diaphysis and beginning of metaphysis.

Biomechanical factors:

1. The bone segment must be large enough to accommodate two rings situated between bone and bone transection.
2. Appropriate distance must be 4-6 cm from joint.
3. Corticotomy must not interfere the joint motion.

Physiological factors:

1. Preservation of Periosteum - which is responsible for bone growth in width.
2. Preservation of Endosteum - Responsible for bone formation and bone resorption.
3. Preservation of local blood circulation.
4. Status of local tissues i.e scar from previous injury, infection or previous surgery can interfere in true regeneration (Osteogenesis).

5. Cyst, porosis and sclerosis can delay or prevent bone formation.

Common mistakes during corticotomy:

1. Too large skin incision.
2. Poor choice of corticotomy level.
3. Uses of over sized osteotome.
4. Destructive hammering of the cortex.
5. Direct cutting of bone marrow canal.
6. Injury to near by vessels and nerves.
7. Performance of twisting corticotome maneuver before complete cutting of medial and lateral cortex.
8. Incomplete corticotomy.
9. Traumatic corticotomy.

Limb lengthening:

Limb length discrepancy (LLD) is a common orthopaedic problem in Bangladesh which arising from either shortening or lengthening of one or more bones in the limb. Poliomyelitis is the most common cause of the LLD, the second important cause is the growth arrest due to osteomyelitis or trauma. Limb lengthening is a long procedure which is associated with many complications but Ilizarov method has least complications which are predictable, preventable and tractable.

Biology:

Distraction osteogenesis

It is a mechanical induction between bony surfaces that are gradually pulled apart in a controlled manner.

1. Gradual lengthening 1mm/day.
2. Faster lengthening leads to failure of bone formation
3. Slower lengthening required in some, can lead to premature consolidation.
4. Rhythm of distraction: increased frequency better e.g. $1/4$ mm 4x/day better than 1mm once a day.
5. Bone formation is in line with direction of distraction.
6. Collagen fibres lined up with direction of distraction. If instability of fixation present collagen fibres become sinusoidal.
7. Mechanism of bone formation is INTRAMEMBRANEOUS.
8. Endochondral bone formation less common but does occur.
9. Fibrous interzone is the layer between the forming columns of new bone.
10. Trabeculae looks like stalactites and stalagmites.
11. Interzone has the undifferentiated mesenchymal cells that can form bone, cartilage or fibrous tissue.

Distraction Histogenesis of Soft Tissues:

1. Gradual distraction is important for soft tissues.
2. Soft tissues lengthening is a combination of stretch and regeneration.
3. Muscle regeneration secondary to addition of sarcomeres as well as recruitment of satellite cells.
4. Nerve regeneration includes new Schwann cells.

Distraction is indicated for:

1. Limb lengthening.
2. Correction of deformities.
3. Bone fragment transport.
4. Stimulus for non-union and Pseudoarthrosis.
5. Neovascularization.
6. Correction of joint contractures.

Complications of limb lengthening:**Bone complications**

1. Premature consolidation.
2. Delayed consolidation.
3. Non union.
4. Axial deviation (secondary deformity)
5. Fracture.
6. Infection.

Joint complications

1. Stiffness
2. Contracture
3. Subluxation
4. Dislocation
5. Arthritis


Soft tissue complications

1. Nerve injury
2. Vascular injury
3. Oedema
4. DVT
5. Muscle contracture
6. Muscle scarring
7. Skin scars

There are many ways to prevent complications. Since distraction related complication occurs gradually they can be identified early and treated early. So frequent follow up (at least every two weeks during distraction and every month during consolidation) are required.

Post operative correction and management:

Before starting deformity correction mathematical calculation has to be done for soft tissue distraction. The equinus is corrected by tightening the anterior threaded rods and loosening of posterior threaded rods. Mean latency period is the 7th day. Correction can be started on the 6th day also. The process of loosening and tightening of connecting threaded rods should be performed 4 times a day @ 0.25mm at 8AM, 12PM, 4PM and 8PM. At the time of correction all hinges should be loosened and after correction all hinges should be tightened if we don't do it the frame will be unstable and regeneration of Achilles tendon may be hampered; but if the tenotomy of Achilles tendon is not done and only soft tissue distraction is done, then hinges remain fixed with nuts, which provides controlled smooth motion. The frame should be retained for a period of 6-8 weeks after full correction or equal to the time needed in achieving the correction except for bone lengthening. Sometimes plaster immobilization is needed after dismantling the frame.



Chapter X

Hormonal Changes in Limb Lengthening Process

Hormonal Changes in Limb Lengthening Process

The synthesis of hormones, products of high biological activity is strictly regulated by actual needs of a macro-organism.

Osteosynthesis with the Ilizarov apparatus in closed fractures is a perfect model for studying of the bone regeneration process in reply to the frame. It has an advantage regarding other methods of treatment: skeletal traction and plaster cast do not provide bone fragment immobilization because traumatic factor is constantly preserved. The osteosynthesis with the Ilizarov lacks all these draw-backs, besides the functional activity of a patient is considerably preserved. In Kurgan the scientific study revealed that a concentration of corticotropin, aldosterone, hydrocortisone, PTH increases right after trauma (3-5 hours). This is a hormonal reply of organism to the trauma. The maximum concentration of those hormones in transosseous osteosynthesis with the Ilizarov apparatus is observed on the 35th posttraumatic day. On the 14th day after trauma, contents of STH and CT increases in blood. On the 5th postoperative day a concentration of all the hormones increased: STH by 3-4 times; CT 2 times, PTH 6 times, adenosine- monophosphate (AMP) 11 times and cyclic guanine monophosphate (cGMP) 6 times.


By the 7th day of distraction the hormonal concentration is reduced by 1-2 times. PTH concentration remains at a high level during 14-30 days of distraction. Calcium (Ca²⁺) and Phosphate concentration is also increased in blood plasma (W.B. High et al 1982).

The most noticeable increase of CT concentration is observed in fixation period; when process of regenerative mineralization is the most intensive. The growth hormone (STH), which concentration exceeds the normal during the whole distraction period, stimulates the synthesis of proteo-glycans and collagen, that is conducive to bone growth. Growth hormone (STH) concentration is 2-3 times higher in a growing child in comparison to healthy adult. (M I Balabolkin, 1978).

In Kurgan, the radiologic and immunologic studies showed that bony tissue regeneration is regulated by hormonal system. Human organism also reacts to distraction rate changes; it reflects in hormonal dynamics.

In Kurgan, an automatic mode of lengthening is more widely used. In comparison to the classic distraction method. When a daily lengthening rate (1mm) is performed in day time by 4 steps (0.25mm X 4 times), the automatic distraction is less traumatic because the lengthening of 1mm is achieved within 24 hours. (-0.017 X 60 times).

Lengthening of a long bone, corticotomy through a sub cutaneous approach, leads to biologically active site. Local undifferentiated mesenchymal cells evolve into osteoblasts producing collagen, osteoid matrix and then bone unireval. Distraction force helps to produce osteogenesis longitudinally in the direction of line of force. The internal distractor, endochondral proliferation, stretches periosteal vessels, nerves, muscle and skin to induce growth. Distraction osteogenesis is the mechanical induction between bony surfaces that are gradually pulled apart in a controlled manner. Distraction osteogenesis is the primary method of bone lengthening.



Chapter XI

Blood Supply and Metabolic Activities of Bony Tissue

Blood Supply and Metabolic Activities of Bony Tissue


In Kurgan (RISC, RTO), experimental studies proved that distraction osteosynthesis stimulates development of dense capillary network. The rate of distraction leads to sufficient increase of volume speed of tibial blood flow by 3 times.

Radio-nuclear studies give the authentic idea about blood flow changes in the extremity being lengthened and metabolic changes constantly connected with it in bony tissue. Metabolic process of both internal and organic bone components constantly and intensively takes place in bony tissue. At present radio-nuclear diagnostics is widely used in studying of metabolic processes in different organs and in bone as well, in diagnostics of different diseases. New osteotropic radio-pharmaceuticals preparations (RPP) e.g. nucleids participating in mineral metabolism is conducive to it.

The activity of metabolic processes in the extremity grows after osteotomy; marked phosphine accumulation in regenerate increased to $452 \pm 0.77\%$ at $P < 0.05$, speed of the volume blood flow to $250 \pm 0.18\%$ at $P < 0.05$.


The capillary blood was sped up in the whole extremity practically within the 1st month of distraction in simultaneous femoral and tibial lengthening. RPP circulation made up to 73% in tibia and 279% in femur. But already by the 3rd month of distraction its circulation increased mostly in femur ($353 \pm 0.23\%$) than in tibia ($302 \pm 0.32\%$).

During the 1st month of fixation the blood circulation practically remains at the same level (350 & 303%). RPP circulation level decreases very slowly. By the 6th month after distraction it is reduced to 24% in femur and to 20% in tibia. Blood circulation normalizes in 12 months after the removal of the apparatus. Intensity of metabolic processes in segments being lengthened changes along with the blood circulation of the extremity.



Chapter XII

Biochemical Procedure of Metabolism in Lower Limb Lengthening in Adult Group



Biochemical Procedure of Metabolism in Lower Limb Lengthening in Adult Group

In Kurgan (RISC, RTO) proteins (albumin, total protein), low-molecular nitrogen containing substances (creatinine, urea, uric acid), lipids (triglycerides and cholesterol), hexoses (glucose, hexosamines), enzymes, alanine amino transferase ALAT; aspartatamino transferase-AsAT, lactose hydrogenase- LDG, hydroxybutirates- dehydrogenase; alkanine phosphates); electrolytes (calcium, magnesium, phosphates, chlorides) were measured in blood stream. Total contents of hydroxy peptide, calcium, magnesium, low-organic phosphates, total 17-oxycorticosteroids were measured in urine. Biochemical tests were performed in patients before treatments, after corticotomy, after lengthening of 1,2,3,4,6,8 cm, in every 30 days of fixation and within one week after removal of apparatus.

During distraction and consequent fixation of the extremity no authentic changes of total protein, albumin, urea, uric acid, glucose, cholesterol, triglycerides, magnesium, chlorides concentration were found in blood serum of the patients. Excretion of total and -amino nitrogen and magnesium did not change in daily urine.

The greatest changes are found in alkaline phosphatase 10 days after distraction its activity increase, 2 time approximately.

* AP is a marker of osteoblast bone synthesis activity and it enhances bone tissue regeneration.

Enhancement of Regenerate Bone Healing

Various biological and mechanical strategies are responsible for improving the ability to enhance the rate and volume of regenerate formation. These are:

- a. Latency period
- b. The performance of low energy osteotomy
- c. Soft tissue preservation
- d. Load sharing fixator designs

Distraction osteogenesis can be characterized into three groups:

1. Failure of adequate regenerate formation in an expected time frame
2. Fracture through the regenerate or adjacent bone
3. Bending of the regenerate after the removal of the frame.

Motorized distraction


8-step distraction (0.125mm increments) compared to 4-steps (0.125mm increments) daily distraction is better. BMD is significantly higher in the 8 step groups than 4-step groups. Motorized distraction may increase patient compliance and comfort.

HBO (Hyperbaric Oxygen Therapy)

HBO increases hypertrophy of cartilage and increases bone formation. In Kurgan center the clinical trial showed that bone healing is increased by HBO and they are routinely using HBO.

Anticatabolic therapy

Biphosphonates: Modern nitrogen containing Biphosphonates (N-bps) prevent fractures from occurring in osteoporotic conditions, control pain. The rationale for the use of N-bps in distraction osteogenesis is to prevent catabolic effects. It increases BMC (bone mineral contents) and BMD and bone volume both in and around the regenerate area.



Chapter XIII

Mechanical Principle of Ilizarov Method

Mechanical Principle of Ilizarov Method

Pearls of Frame Mounting

Rule of Two's

1. 2 cm between skin and frame.
2. 2 rings/bone segment
3. 2 points fixation/ring
4. $2 \times 2 = 4$ connecting rods between rings
5. Fixation both 2 ends of the bone segment (near-near and far-far)
6. Pin and wire fixation in 2 planes.

Methods to Increase Frame Stability:

Rings	Decreasing ring diameter Decreasing ring to skin distance Securing near and far ends of bone segment Increasing number of rings (use "dummy ring" to span long distances)
Ring connections	Increasing number of connections Increasing rigidity of connections (telescopic rods to span long distances)
Wires	Increasing number of wires Increasing diameter of wires Maximizing crossing angle wires Opposing olive wires Drop wires
Bone considerations	Maximize bone end contact Apply compression / distraction

N.B.: Don't violate the original Russian Rule of Technique.

What Not To Do?

Some of the following cases were done outside Bangladesh and in Bangladesh.

Case No. 1



Picture 1: Ring sizes are much more bigger, rings that are too large do not support the transfixing wires adequately and osteogenesis impaired. No rubber stopper, no dressing gauge is seen.



Picture 2: Close up view of the rings. Violation of mechanical stability (lose of trampoline effect), 4 cm LLD persists.

Cont...

Case No. 2



Picture 3: 2 fingers in the lateral side, 5 fingers in the middle side.



Picture 4: 5 fingers in the medial side, 2 fingers in the lateral side.



Picture 1: 3 fingers in the lateral side and one finger in the medial side.



Picture 2: Violation of mechanical stability, no dressing gauge is seen. 4 fingers in the medial side and one finger in the lateral side.

Case No. 3



Picture 5: Close up view, ugly frame assembly.



Picture 1: No proper ring sizes. No mechanical stability, No rubber stopper, No dressing gauge is visible.

Case No. 3



Picture 2: Close up view of the frame, Wires are loose, No rubber stopper and no dressing gauge is visible.



Picture 3: 3 fingers in the lateral side and one finger in the medial side.



Picture 4: No dressing is seen, no rubber stopper, foot is equinus position.

Case No. 4



Picture 1: Radiograph of right lower tibia fibula fracture with 2 rings in situ which is not adequate in case of adult patient.



Picture 2: Two rings are not adequate. Rings are not in proper position, mechanically unstable.



Picture 3: 4 fingers in the medial side and 1 finger in the lateral side, This is the violation of the rule.

Chapter XIV

Different Plane (Frontal plane, Sagittal plane and Horizontal plane) Deformities and Bio-mechanics of the Ankle and Foot

Different Plane (Frontal plane, Sagittal plane and Horizontal plane) Deformities and Biomechanics of the Ankle and Foot

Frontal/Coronal Plane Deformities of Ankle and Foot

Clinically frontal plane is best evaluated from behind. In the frontal plane the axis of the body of the calcaneus is normally parallel to the anatomic axis of the tibia. Heel varus and heel valgus are seen in this plane and the plane of deformity is hind foot.



Figure 14.1a: Varus heel.



Figure 14.1b: Valgus heel.

Sagittal Plane Deformities of Ankle and Foot

This plane is visible when we see the deformity from the side. Pes equinus, Pes Equinocavus, Pes planus and Pes calcaneus are seen in this plane and the place of deformities we see in hind foot, mid foot and forefoot (Figure 14.2a-14.2e).



Figure 14.2a: Pes Equinus.



Figure 14.2b: Pes Equinocavus.



Figure 14.2c: Valgus heel.



Figure 14.2d: Pes Planus.



Figure 14.2e: Pes Calcaneus.



Figure 14.3c: Planes of ankle and foot (Horizontal, sagittal and frontal).

Horizontal plane deformities of ankle and foot:

This plane is visualized when we see the deformity from the up-down. Adduction & abduction of the fore foot are seen in this plane.



Figure 14.3a: Adducted foot.



Figure 14.3b: Pes Planus.

The human foot is complex structure adopted to allow orthograde bipedal stance and locomotion. It is the only part of the human body which is always in contact with the ground. There are 28 major bones in the foot, 31 major joints including the ankle joint. Functionally the skeleton of the foot may be divided into tarsus, metatarsus and phalanges.

Arches of the Foot

Three main arches are recognized in the foot.

Medial longitudinal arch.

It is made of the calcaneus, talus, navicular and three cuneiform and three metatarsals. The pillars are the posterior aspect of the calcaneus and three metatarsal heads.

Lateral longitudinal arch:

The bone making up the longitudinal arch are the calcaneus, the cuboid and the 4th & 5th metatarsals. The pillars are calcaneus and the lateral 2 metatarsal heads.

Transverse arch:

The bones involved here are the bases of the 5 metatarsals, the cuboid and the cuneiforms.

Biomechanics of ankle and foot:

Planes of motion: Plantar flexion and dorsiflexion refers to movement in the sagittal plane and occur principally; but not exclusively at the ankle, metatarsophalangeal and interphalangeal joints.

Inversion is tilting of the plantar surface of the foot towards the midline.

Eversion is tilting away from the midline, this is motion in coronal/ frontal plane.

Different planes and deformities:

Planes and Deformity	Place of Deformity	View
1. Sagittal Plane		
a) Equinus	Ankle	To be Seen from side
b) Equinovarus	Hind foot,	
c) Pes planus	Mid foot	
d) Pes planovalgus	and	
e) Pes calcaneus	Fore foot	
2. Frontal Plane		
a) Heel varus	Hind foot	To be seen from back
b) Heel valgus	deformity	
3. Horizontal Plane		
a) Fore foot adduction	Fore foot deformity	To be seen from up to down
b) Fore foot abduction		

Adduction is the movement of the foot towards the midline in the transverse plane.

Abduction is the movement away from the midline. This movement occurs at the transverse tarsal joints and to a limited degree, the first tarso metatarsal joint and the metatarsophalangeal joints.


Supination describes a three dimensional movement and is a combination of adduction, inversion & plantar flexion.

Pronation is the opposite motion i.e., a combination of abduction, eversion and dorsiflexion.

Humans are bipedal. We are plantigrade, i.e., we set the whole length of the foot down on the ground; whereas most mammals are digitigrade i.e., they stand and walk on their toes. Mid foot is responsible for dynamic distribution of weight through medial longitudinal arch.

Weight is transmitted from hind foot to fore foot through the transverse arches.

Any abnormality from the plantigrade will lead to deformities in the foot (Table 14.1).



Chapter XV

Disorders of the Bone Configuration and Function of Leg Bones

Disorders of the Bone Configuration and Function of Leg Bones

I. Deformities of leg bones

II. & III. Pseudoarthroses of leg bones and Defects of Leg Bones

IV. & V. Leg Lengthening, Leg shortening with segmental axis disturbance leg shortening without its axis disturbance



I. Deformities of leg bones

Acquired Deformities of Leg Bones:

- Post traumatic (after multiple polyfragmental fractures - 42.1% of all the injuries)
- Post infection (sequelae of haematogenous osteomyelitis - 71.2%)
- Flaccid pareses and paralyses (60-75%).
- Metabolic and dystrophic (sequelae of rickets, rickets-like diseases, osteoarthritis).

Congenital Deformities of Leg Bones (Dysplastic Systemic Skeletal Diseases)

- Dyschondroplasia (Ollier's disease)
- Blaunt's disease
- Osteogenesis imperfect
- Fibrous dysplasia
- Developmental anomalies

Classification of Leg Bone Deformities According to Planes

- Uniplanar
- Biplanar
- Multiplanar

Classification of Leg Bone Deformities by Localization

- Metaphyseal
- Diaphyseal
- Metadiaphyseal (subcondylar)
- One-level (Monofocal)
- Multilevel (Polyfocal)
- Longitudinal
- Transverse

Classification of Leg Bone Deformities by The Number of Segments

- Monosegmental
- Polysegmental

Sequelae of Haematogenous Osteomyelitis are Characterized By

- Multicomponent deformity of leg bones in the proximal part (52.9%)
- Multicomponent deformity of leg bones in the distal part (17.7%)
- Varus deformity of leg bones in the proximal part (11.8%)
- Varus deformity of leg bones in the distal part (5.9%)
- Outward and inward torsion

- Shortening of limb segments (in all patients)
- The knee instability (57.7%).


Dyschondroplasia (Ollier's Disease) is Characterized by

- 1-25 cm shortening of the lower limb (by an average of 9.8 cm)
- Preferential deformity localization in proximal leg (63.1%)
- The presence of the knee and ankle contractures (45%)
- The knee instability (75%).


Stages of Preoperative Planning

- Determination of true deformity plane (CORA)
- Calculation of the amount of deformity correction and lengthening
- Biomechanical designing
- Determination of osteotomy levels
- Determination of osteotomy shape
- Selection of deformity correction type


A diagram of hinge placement at the apex of deformity (Figure 15.1).



The amount of the base of wedge-shaped regenerated bone



The amount of bone, for formation of trapezoidal regenerated bone




Stages of Surgery

- Osteosynthesis of leg bones (insertion of wires, mounting of the Ilizarov fixator, placement of hinge units for deformity correction)
- Skin incision, approach to bone
- Osteotomies
- Deformity correction (partial or complete -up to 150)
- Skin closing
- Control x-rays

Main Principles of Transosseous Osteosynthesis for Deformity Correction

- Insertion of wires with stoppers in the process of deformity correction
- Hypercorrection of the external supports of the transosseous fixator during its mounting
- Proper and efficient arrangement of the fixator hinge units

A variant of the fixator configuration for correction of varus deformity of the leg upper third with the knee «protection» (Figure 15.4 & 15.5).



A variant of the fixator configuration for correction of valgus deformity of the leg upper third.



Stages of Postoperative Management

- Gradual correction of residual deformity
- Proper dressings
- Adequate functional and static weight bearing of the limb
- Efficient exercise therapy
- Controlling the regenerated bone condition

Periods of the fixator removal are determined by the followings

- The mean periods of consolidation depending on the amount of deformity, limb shortening, patient's age, etiology
- The presence of the X-ray signs of newly formed bone
- The data of clinical testing the consolidation stability

Principles of removing the transosseous ring fixator

- Adherence to the indications for the fixator removal (consolidation, clinical signs and those by x-rays),
- Sound anaesthesia
- Decreasing the forces in «fixator-bone» system
- Adherence to asepsis and antisepsis

A patient should observe the following rules after the removal of Ilizarov fixator

- Axial and functional loading of the limb should be gradually increased
- Adequate exercise therapy should continue
- Regular procedures of physical therapy
- Control of subsequent examinations

Tactical errors (Calculated errors)

- The wrong selection of the number and levels of osteotomies
- The faults in the calculations of limb segmental deformity and shortening
- Neglecting the disease etiology and patient's age

Technical errors

- In the preoperative period- the wrong selection of the Ilizarov fixator supports and parts
- While performing surgeries - non-observing the rules of wire insertion, biomechanical principles of mounting the fixator supports and units, making osteotomies, neglecting the creation of soft tissue reserve during wire insertion and fixation

The errors that we should keep in mind in postoperative period

- Unreasonable frequently changing the fixator
- Non-observance of deformity correction rates
- The absence of timely x-ray control of the dynamics of bone regeneration
- Incorrect interpretation of x-rays
- Premature removal of the fixator
- Underestimation of the importance of exercise therapy and the possibilities of early limb weight-bearing with the fixator applied

Complications


- Inflammation of soft tissues at the site of wire skin interface.
- Postoperative neuropathy of the peroneal nerve
- Consolidation in osteotomy site
- Cutting wires out of bone
- Equinus foot deformity
- The knee contracture
- Pseudoarthrosis formation
- Subluxation of the joint
- Transformation of the regenerated bone

II & III. Pseudoarthroses and Defects of Leg bones

Pseudoarthrosis is a nonunion of bone in the average statistical period of time. True bone defects, means any loss of bone substance.

The true bone defect represents the total amount of interfragmental diastasis and anatomic segmental shortening. This takes into account the volume and amount of expected bone tissue loss in process of the planned removal of nonviable grafts, resection of the ends of fragments and their required the duplication at the docking site (Figure 15.6-15.9) and Table 15.1.


Classification of defects and pseudoarthroses



Defect-pseudoarthrosis

Defect-Diastasis

Classification by the amount of defect



Total

subtotal

partial

Classification by the anatomic and radiographic criteria of fibula condition and soft tissue retraction of leg



1

2

3

4

5

6

1- Preserved fibular integrity;

2- Fibular dislocation (subluxation) in the distal syndesmosis;

3- Fibular dislocation (subluxation) in the proximal syndesmosis;

4- Fibular defect-diastasis;

5- Fibular overlapping;



6- The presence of tibiofibular synostosis Etiolog

Pseudoarthrosis characterized by the type of articular ends (callus formation)

Defect (pseudoarthrosis Type)	Mobility at the Defect (Pseudo-arthrosis) Site	description of Fragment Ends	The Data of Radio include and Arteriographic Study
Hypoplastic	Excessive	Atrophy, sharpening, absence of periosteal growth, medullary canal eburnation	Accumulation of labelled pyrophosphate at the fragment ends is close to the norm, blood flow is decrease. Vascular network is poor.
Normoplastic	Moderate	Not changed or moderately osteoporotic, end-plates look like narrow sclerotic strips	Accumulation of labeled pyrophosphate at the fragment ends is 1.5-2=fold increased. blood flow is moderately accelerated. Hyper vascular.
Hyperplastic	Stiff	"Bamboo"-shaped thickening, the interfragmental gap is twisted and interrupted, end-plates are marked but their boundaries are unclear	Significant accumulation of marked pyrophosphate. Circulation in the interfragmental gap is 3-fold accelerated. Marked formation of vascular collaterals in the soft tissues surrounding the zone of nonunion.
Neoarthrosis		The surface of fragments is smooth, convex and concave, an "articular" gap is present, medullary canals are closed by bone tissue over considerable extent.	Marked hypervascularization of soft tissues.

Classification by the shape of fragment ends:

Diagrams of the «articular» ends of pseudoarthrosis zone



1, 2, 5- congruent;
3, 4-incongruent;
1- transverse;

2, 3-oblique;
4-oblique-and-transverse;
5- hinged Congenital

Etiolog:

- Congenital
- Acquired
 - a) Posttraumatic;
 - b) Post infective;
 - c) Postsurgical.

With regards to infection

- Complicated by pyogenic infection;
- Uncomplicated by pyogenic infection

Osteogenesis in pseudoarthrosis zone

- Hypertrophic
- Normotrophic
- Atrophic.

Aim of Treatment and Rehabilitation Process - Recovery of Limb Anatomic-and-Functional Integrity

- Bone defect filling
- Integrity recovery of segmental bone
- Segment length restoration
- Correction of segmental deformities
- Correction of the defective contractures of adjacent joints

Main Approaches in The System of Reconstructive-And-Rehabilitative Treatment of Patients with Leg Defects by Ilizarov Technique

- Union at the docking site of bone fragments
- Lengthening of fragments
- Tibiofibular synostosis
- Closed gradual distraction of interfragmental tissues with formation of regenerated bones.

Types of Monofocal Transosseous Osteosynthesis

- Monofocal combined distraction-compression osteosynthesis
- Monofocal distraction-sequential compression osteosynthesis
- Monofocal alternate compression-distraction osteosynthesis.

Indications for Monofocal Osteosynthesis

The presence of tibial pseudoarthrosis, including that with fixed angular deformity and anatomic segmental shortening not exceeding 2 cm.

Indications for Bifocal Osteosynthesis

The presence of true tibial defect > 2-3 cm with anatomic segmental shortening and without it.


Types of bifocal transosseous osteosynthesis

- Bifocal simultaneous compression-distraction osteosynthesis with fragmental lengthening
- Bifocal sequential compression-distraction osteosynthesis with fragmental lengthening (Figure 15.10).

A variant of the Ilizarov fixator configuration for bifocal transosseous osteosynthesis





Principles of Transosseous Osteosynthesis for Pseudoarthrosis Compression of bone fragments along bone axis



Side-to-side compression of bone fragments using posts

Side-to-side compression of bone fragments using olive wires

Filling of extensive bone defects



Tibiofibular Synostosis and Tibialization

Techniques of tibiofibular Synostosis

- Using a split-fragment;
- Using a cylinder-fragment

Indications for tibiofibular Synostosis

The presence of total (subtotal) tibial defect, when defect filling by fragmental lengthening is not possible. It is very effective in young person's only.


Classification of Tibiofibular Synostosis

Techniques by Technological Features

- Bypass tibiofibular Synostosis
- Fibula Tibialization (Figure 15.14 & 15.15).



Indications for Polyfocal Osteosynthesis:

The presence of true tibial defect > 2-3 cm with anatomic segmental shortening and without it



Fragmental multilevel lengthening

Combination of techniques



Complications of the Following Structural Masses Can be Observed

- Segmental soft-tissue components (skin, subcutaneous fat, muscles, vessels, nerves);
- Adjacent joints and articulations of twin bones;
- Bone fragments and regenerated bone.

Complications of Skin and Subcutaneous Fat

- Inflammation in the zone of contact with inserted wires.
- Soft tissue suppuration in places of compression produced by the fixator parts in case of Their improper selection.
- Local limited erosive wound surfaces around the exit of wires may be observed.
- Dermatitis.

Complications of Muscles

- Insertion of wires through muscular masses during osteosynthesis may result in limiting some motor activity of the limb involved.
- Complications of vessels and nerves.
- Bleeding at the site of wire exit.
- Erosive arterial bleeding.
- Oedema.
- Injury of nerve trunks.
- Failure of innervation and limb segment being lengthened.

Complications of Adjacent Joints and Twin Bone Articulations

- Purulent arthritis
- Temporary or persistent limitations of movements in the neighboring joints
- Development of subluxation.

Complications of Bone Fragments and Regenerated Bone

- «Wire» osteomyelitis-cortical osteomyelitis
- Osteomyelitic process aggravation
- Fracture at the level of wires
- «Cutting out» of wires from the soft tissues
- Transverse, angular or peripheral displacement of bone fragments
- Formation of angular deformities in the zone of docking bone fragments or regenerated bone.



IV & V. Leg Lengthening, Leg shortening with segmental axis disturbance leg shortening without its axis disturbance

Generally Monofocal distraction osteosynthesis of the leg is indicated for segmental lengthening up to 5 cm Bifocal distraction osteosynthesis of the leg is indicated for segmental lengthening above 5 cm (Figure 15.17).

Possible Complications During Leg Lengthening

- Inflammation of soft tissues around the wires - 18%
- Pareses development during distraction - 5.6%
- Development of the knee and the ankle contractures, foot deformities- 60%
- Regenerated bone fracture after the fixator removal - 3.2%

Variants of the Ilizarov fixator configurations for bifocal lengthening of the leg



Case Study 1

Reconstruction of the Limb Versus Amputation

1. Open degloving injury of left leg with bone loss of left tibia GIIIB.
2. Radiograph of left tibia fibula with bone loss in the lower tibia.
3. Close up view of the left leg with Ilizarov in situ.
4. Ilizarov in the left leg after 1 month follow up.
5. Close up view of the left leg with Ilizarov in situ.
6. Skin grafting was done.
7. After 2 month follow up
8. Radiograph of tight tibia fibula with distraction osteogenesis.
9. After 3 months follow up.
10. Radiograph of left tibia fibula with good regenerate in the upper tibia .
11. After 4 months follow up.
12. After 4 ½ months follow up.Front view.
13. After 4 ½ months follow up.back view.
14. After 6 months follow up.
15. After 6 months follow up
16. n OR Table before removal of Ilizarov apparatus.
17. Radiographic result after 6 months.
18. Clinical appearance of the patient.



Cont....



Case Study 2

1. Open comminuted fracture left tibia G IIIA
2. Radiograph of comminuted fracture of left tibia.
3. External view of left leg and ankle with Ilizarov in situ.
4. Skin grafting was done.
5. 47 years old lady with Ilizarov apparatus, 2 months after the application of Ilizarov.
6. Healing of tissues with Ilizarov apparatus.
7. External view of the leg and ankle.
8. Radiograph of tibia fibula with Ilizarov in situ.
9. After 4 months follow up: correction of toes also visible.
10. Radiographic final result.



Case Study 3

1. Crush injury of right lower femur with right tibia fibula and ankle GIIIB (Floating knee)
2. 14 years old girl with Crush injury of right lower femur with right tibia fibula and ankle GIIIB
3. Exposed right whole tibia with Ilizarov in situ, patient in OR
4. After 1 month follow up the right leg with Ilizarov fixator
5. After 1 ½ follow up, right leg with Ilizarov fixator
6. Almost covered the whole tibia with granulation tissue
7. The whole tibia is covered by a good granulation tissue
8. Skin grafting was done
9. Radiographic view of tibialization with Ilizarov in situ
10. The whole leg is covered by skin and bone is stabilized by Ilizarov ring fixator
11. Smiling patient after 5 ½ months follow up
12. Clinical appearance of the patient after 7 months



Case Study 4


Open Fracture Tibia and Fibula

1. Open comminuted fracture right tibia fibula GIIIA
2. Radiograph of right tibia fibula
3. Ilizarov fixator in the right tibia fibula, 6th post OP, 24 years old male
4. After 21 days follow up
5. Patient can walk with Ilizarov
6. Radiographic final result after 4 months
7. Clinical appearance of the patient after 4 months



Mal Uniting Fracture Right Lower Tibia Fibula (3 & ½ months old) GII

1. 27 years old male with mal uniting fracture of right lower tibia fibula (3 & ½ months old) with varus deformity (front view).
2. 27 years old male with mal uniting fracture of right lower tibia fibula (3 & ½ months old) with varus deformity (Back view).
3. Radiographic view of right tibia fibula with maluniting fracture.
4. Radiographic view of after the application of Ilizarov apparatus, 2nd post op.
5. Radiographic view of almost corrected varus deformity with union before removal of Ilizarov apparatus, 5 & ½ months follow up.
6. External view of the Ilizarov apparatus in the right leg during treatment.
7. Final Radiographic result of right tibia fibula after removal of Ilizarov apparatus.
8. Smiling patient in sitting position after 7 months follow up.
9. Clinical appearance of the patient after 8 months follow up.



Case Study 6


Open Fracture Right Lower Tibia Fibula G-IIIa

1. Radiograph of right lower tibia fibula with bone loss, uniaxial fixator in situ.
2. 48 years old male during treatment with Ilizarov apparatus.
3. Radiograph of right tibia fibula with Ilizarov fixator in situ. Docking and corticotomy site is clearly visible.
4. After removal of the fixator, 6 months follow up.
5. Radiograph of right tibia fibula after 8 months follow up, good union is achieved.
6. Clinical appearance of the patient in sitting posing.
7. Clinical appearance of the patient in standing position



Infected non Union Right Tibia Fibula with Interlocking Nail in situ GIIIB, Nail is Exposed


1. 22 years old male, Infected non union right tibiafibula with interlocking nail in situ GIIIB, nail is exposed.
2. Close up view of right upper tibia fibula, exposed nail with bone.
3. Radiograph of right tibia fibula, interlocking nail with non union.
4. Radiograph of right tibia fibula with gap afterremoval of inter locking nail.
5. Radiograph of right tibia fibula with segmentalbone transport(arrow marking), Ilizarov in situ.
6. Radiograph of right tibia fibula with segmentalbone transport(arrow marking), with good regeneration and consolidation after 5 months follow up, with Ilizarov in situ.
7. Clinical appearance of the patient with Ilizarov apparatus in the leg.
8. Smiling patient with healing of the exposed bone.
9. Final Radiographic result after 7 months follow up.
10. Clinical appearance of the patient after 7 months follow up.



Case Study 8

Infected Open Fracture Right Tibia Fibula GIIIB with Exposed Plate & Screws

1. & 2. Infected open fracture right tibia fibula GIIIB with exposed plate & screws
3. After removal of the plate, screws & dead bone and debris
4. 45 years old male with Ilizarov fixator in the right leg
5. Radiographic view right tibia fibula with Ilizarov fixator in situ, after 2nd post OP
6. Radiographic view of right tibia fibula with Ilizarov fixator in situ, after 1 month follow up.
7. Radiographic view of right tibia fibula with Ilizarov fixator in situ, after 3 months follow up.
8. Radiographic view of right tibia fibula with Ilizarov fixator in situ, after 9 months follow up.
9. Patient is in standing position with Ilizarov apparatus in the right leg
10. Final radiographic result after 10 months follow up.
11. Clinical appearance of the patient with smiling face after 11 months follow up.



Open Fracture (GIIIa) Left Tibia Fibula

1. Open fracture (GIIIa) with uniaxial Ex-fix in situ.
2. 23 years old girl, open fracture left tibia fibula (GIIIa); uniaxial Ex-fix in situ
3. Radiographic view of left tibia fibula with uniaxial Ex-fix in situ.
4. In OR with Ilizarov apparatus.
5. Clinical appearance of the patient after removal of Ilizarov fixator.
6. Radiographic consolidation of left tibia fibula after 6 months follow up.
7. Clinical appearance of the patient after 3 years follow up.
8. Radiographic result of left tibia fibula after 3 years follow up.



Case Study 10

Open Comminuted Fracture Left Lower Tibia Fibula (GII)

1. Radiograph of left lower tibia, open comminuted fracture (G II).
2. 50 years old male – 2 weeks after surgery with Ilizarov device in the left leg and ankle.
3. 50 years old male – Ilizarov in situ, after 1 month.
4. Radiograph of left tibia and ankle, healing is going on, after 3 months follow up.
5. Radiograph of left tibia and ankle - 5 months follow up after removal of fixator.
6. Clinical appearance of the patient, after 9 months follow up.
7. Patient is in sitting position, after 9 months follow up.



Comminuted Fracture Right Lower Tibia Fibula

1. 52 yrs. old male comminuted fracture right lower tibia and fibula.
2. Radiographic view of right lower tibia and fibula with Ilizarov in situ.
3. Clinical appearance of the patient with Ilizarov apparatus in the right lower leg.
4. Radiograph of right lower tibia fibula after 7 months follow up.
5. Clinical appearance of the patient in sitting position after 7 months.
6. Clinical appearance of the patient after 1 year standing position.



Case Study 12

Open Segmental Comminuted Fracture Left Lower Tibia GII

1. 42 yrs old male, Open segmental comminuted fracture of left lower tibia G II. Radiographic view of right lower tibia and fibula with Ilizarov in situ.
2. Radiographic view of left tibia fibula with Ilizarov in situ; 2nd post op. Radiograph of right lower tibia fibula after 7 months follow up.
3. Radiographic view of left tibia fibula with lengthening of fibula is seen.
4. In OR before dismantling the Ilizarov apparatus.
5. Radiographic final view of tibia fibula after 9 months.
6. Clinical appearance of the patient after 9 months (Sitting position, standing front view & back view).



Non Union Left Upper Tibia

1. 8 years old girl – Deformity of left knee, heel is facing forward.
2. Radiographic non union left upper tibia with deformity of fibula and shortening (5 cm).
3. Side view of the leg.
4. Radiographic result of left tibia fibula with Ilizarov fixator in situ.
5. During treatment with the Ilizarov fixator.
6. Smiling patient with the Ilizarov fixator.
7. Radiographic result of left tibia fibula.
8. Clinical appearance of the patient after full correction.



Case Study 14



Aseptic Nonunion of Long Bones with Anterior Bowing

1. Radiograph of Non union of right tibia with anteromedial bowing after implant failure.
2. Radiograph of right tibia fibula with Ilizarov in situ after 2 months follow up
3. Radiograph of right tibia fibula with Ilizarov in situ after 4 months follow up
4. Radiograph of right tibia fibula with full consolidation and deformity correction
5. Clinical appearance of the patient after 6 months follow up.
6. Clinical appearance of the patient after 6 months follow up
7. Patient is in standing position.
8. Patient can squat easily



Infected Gap non Union Right Lower Tibia Fibula, Equinus Foot with AO Fixator *in situ*, Treated in Apollo Hospital, Dhaka

1. 36 years old male, External view of the AO fixator in the right leg and foot, patient in OR (Bari-Illizarov orthopaedic center.)
2. Close up view of the AO fixator in situ.
3. Radiographic view of the right tibia fibula and ankle with AO fixator in situ.
4. Radiographic view of the right tibia fibula with Ilizarov fixator in situ. follow up after 10 days, corticotomy site is visible with 2 guide wires in tibia and fibula.
5. Radiographic view of the right tibia fibula and ankle after 2 months follow up. Corticotomy and docking sites are nicely visible.
6. Smiling patient with Ilizarov fixator in right leg and ankle after 5 months follow up.
7. Radiographic view of the right tibia fibula and ankle, after 6 months follow up.
8. Clinical appearance of the patient, final outcome after 7 months (front view).



Case Study 16

Gap Nonunion of Left Tibia with Gross Deformity and LLD (12.6)

1. & 2. Posterolateral bowing of left leg with 12.6 LLD.
3. Clinical photograph of 14 years old boy before surgery.
4. Radiograph of hypertrophic deformed fibula with gap nonunion of left tibia, before surgery.
5. Radiographic result of distraction osteogenesis with correction of deformity is seen.
6. Patient with Ilizarov apparatus after 8 months follow up.
7. Radiographic result of tibia fibula.
8. & 9. Clinical appearance of the patient after 14 months. No LLD, No deformity.



Non Union Left Femur Due to Gunshot Injury, 13 cm LLD

1. & 2. 40 Yrs. old male. 19 yrs. of non union left femur due to gunshot injury, 13 cm shortening, before surgery.
3. Radiograph of non union of left femur with gunshot injury.
4. 13 cm LLD in lying position.
5. 1st post op with Ilizarov in situ.
6. 5 cm length is gained after 60 days.
7. Monofocal corticotomy of left tibia with fibula radiograph of osteotomy.
8. No more LLD (limb length discrepancy). Ilizarov in situ after 8 months follow up.
9. Radiograph of left tibia fibula with good regenerate.
10. Radiographic union of left femur with gunshot injury.
11. After removal of Ilizarov fixator 10 months follow up. Lengthening of tibia and fibula is visible.
12. Clinical appearance of the patient, front and back view. Limb length equalization is seen.



Case Study 18


Infected Gap non Union Right Lower Tibia, Deformed Foot with Biaxial Fixator *in situ*

1. 35 years old male, Infected gap non-union right lower tibia, deformed foot with biaxial fixator *in situ*.
2. Radiograph of the same patient, tibia and foot with biaxial fixator *in situ*.
3. Close-up view of right leg and foot with biaxial fixator *in situ*.
4. Smiling patient with Ilizarov fixator in leg and foot.
5. Close-up view of Ilizarov fixator in leg and foot after 3 months follow up.
6. Patient with Ilizarov fixator in the leg and foot after 4 months follow up.
7. Radiographic view after the removal of the Ilizarov fixator.
8. External view of the patient leg and foot after 6 months follow up (sitting position).
9. Final follow up of the patient after 8 months, full correction is achieved, patient is in standing position (front view).



Osteomyelitis of Right Upper Tibia Leading to Gap non Union


1. Radiographic view of right upper tibial chronic osteomyelitis.
2. Radiographic view of same right tibia after 2 years leading to gap non union of right upper tibia.
3. 7 years old girl during treatment with Ilizarov apparatus.
4. Radiographic view of right tibia fibula with Ilizarov in situ, distal corticotomy is done.
5. Final radiographic result after 4 months follow up, good union is achieved.
6. Clinical appearance of the patient in sitting positing.
7. Clinical appearance of the patient in standing positing.



Case Study 20

Chronic Osteomyelitis of Left Tibia

1. 11/2 yrs. old baby, Radiographic view of chronic osteomyelitis of left tibia.
2. Radiographic view of left tibia after application of Ilizarov with compression distraction osteosynthesis.
3. Clinical appearance of the patient with Ilizarov apparatus in the left leg.
4. Radiographic final result of left tibia after 4 months.
5. Clinical appearance of the baby with smiling face (sitting position).
6. Clinical appearance of the baby with smiling face (Standing position).



Non Union of Right Lower Tibia with Broken Nail



1. Radiograph of right tibia with broken nail *in situ*
2. Radiograph of right tibia fibula with Ilizarov in situ after 1 month follow up
3. 24 years old male with Ilizarov in the right leg
4. After 2 months follow up
5. Patient can walk with Ilizarov frame easily
6. Radiograph of right tibia with good regenerate
7. Radiographic result with good union
8. Clinical appearance of the patient after 6 months follow up
9. Patient can squat easily



Case Study 22

Post Traumatic Left Tibia Vara

- 1,2 & 3 27 years old male with post traumatic leftsided tibia vara (Bad scar) (Front & Back View)
4. Radiographic view of the left knee before surgery
 5. Radiographic view of the left knee with Ilizarov apparatus, visible opening wedge osteotomy with hinges
 6. External view of the leg with Ilizarov apparatus *in situ*
 7. Final Radiographic result after surgery
 8. Clinical appearance of the patient after 6 years follow up, no tibia vara, discoloration of the skin disappeared



Post Traumatic Tibia Vara with Bad Scar

1. Post traumatic left tibia vara (Front view)
2. Back View
3. Sitting position Galeazzi sign (+)
4. During squatting deformity of left knee is seen.
5. After the application of Ilizarov, 2nd post-op.
6. 14 years old boy can walk with the Ilizarov apparatus.
7. Radiographic final result after 4 months follow up.
8. Clinical appearance of the patient (No Deformity).
9. No deformity is seen when squatting.



Case Study 24

Right Anteromedial Bowing

1. Age 28 years old man - Right Anteromedial bowing, front view.
2. Right Anteromedial bowing, (back view)
3. Radiograph of right tibia (before treatment)
4. During treatment, with Ilizarov.
5. After removal of Ilizarov fixator, with plaster immobilization.
6. The radiographic result.
7. Final follow up clinical appearance of the patient
8. Patient is happy with the Ilizarov treatment.



Tibialization

1. Big gap non union of left tibia with uniaxial fixator *in situ*
2. 27 years old male with Ilizarov in the left leg
3. Patient can walk with the Ilizarov apparatus
4. Radiographic view of tibialization with Ilizarov *in situ*
5. Radiographic result after the removal of Ilizarov apparatus
6. Just after the removal of the Ilizarov apparatus
7. Radiographic result after 6 months follow up



Case Study 26

1. Radiographic view of left tibia with very big gap
2. Same patient with uniaxial fixator in situ with bad scur
3. Tibialization of left tibia with Ilizarov
4. Radiographic view of tibialization, left sided after 5 months
5. Radiographic view of tibialization, left sided after 6 months
6. Radiographic result after removal of the Ilizarov apparatus
7. Clinical appearance of the patient.





Post Osteomyelitic Big Gap non Union of Right Tibia.

1. Post osteomyelitic big gap non union of right tibia, 7 years old boy.
2. Sarpentine deformity of right leg, front view.
3. Sarpentine deformity of right leg, back view.
4. During treatment with Ilizarov apparatus, in OR.
5. Radiograph of tibialization with Ilizarov in situ.
6. Follow up after 2 months with Ilizarov apparatus.
7. Radiograph of tibialization with Ilizarov after 4 months follow up.
8. After dismounting Ilizarov apparatus, 5 months follow up.
9. Clinical appearance of the patient, sitting position.
10. Clinical appearance of the patient 5 cm LLD persists, front view.
11. Clinical appearance of the patient 5 cm LLD persists, back view.
12. Relengthening of the tibialization part after 1 year, radiograph of lengthening part with Ilizarov in situ.
13. Clinical appearance of the patient with Ilizarov in situ after 2 months follow up.
14. Clinical appearance of the patient with Ilizarov in situ after 3 months follow up.
15. Final radiograph of relengthening part of tibialization after 6 months.
16. Clinical appearance of the patient, front view after 6 months.
17. Clinical appearance of the patient, back view after 6 months.



Cont...



Hockey Stick Deformity (C-Type Deformity)

1. Age 18 years old girl – Hockey stick deformity of right ankle with 5 cm shortening.
2. Radiographic view, hockey stick deformity.
3. Right deformed ankle with bad scar.
4. During treatment with Ilizarov ring fixator.
5. During treatment after 2 months.
6. After removal of the Ilizarov apparatus.
7. Deformity is corrected and limb length equalization is achieved.
8. Radiographic result of right ankle deformity.
9. Radiographic view of whole tibia and ankle after correction.



Case Study 29

Anterolateral Bowing

1. Age 9 years old girl– Left Anterolateral bowing with ankle valgus.
2. Age 9 years old girl – Before treatment.
3. Before treatment.
4. Before treatment.
5. Radiograph before treatment.
6. Radiograph before treatment.
7. During treatment.
8. During treatment.
9. During treatment after 4 months.
10. The Radiographic result.
11. The Radiographic result.
12. During treatment.
13. Final follow up.
14. Final follow up.
15. With AFO.
16. Final follow up after 1 year (Back View)
17. Final follow up after 1 year (Front View)
18. Final follow up after 2 years
19. Final follow up after 2 1/2 years



Cont...



Case Study 30

Left Anteromedial Bowing

1. Age 9 years old boy - Left tibia Anterolateral, bowing front view.
2. Left tibia Anterolateral, bowing (back view).
3. Anterolateral, bowing of left tibia.
4. Café au lait.
5. Radiograph of both tibia fibula before treatment.
6. Correction of Deformity with Ilizarov device.
7. The Radiographic result.
8. During treatment with Ilizarov device.
9. During treatment (back view).
10. Correction of the deformity.
11. Correction of the deformity.
12. Final follow up.
13. With AFO.



Rt Anteromedial Bowing

1. Age 28 years old man - Right Anteromedial bowing, front view.
2. Right Anteromedial bowing, (back view)
3. Radiograph of right tibia (before treatment)
4. During treatment, with Ilizarov.
5. After removal of Ilizarov fixator, with plaster immobilization.
6. The radiographic result.
7. Final follow up.
8. Patient is happy with the Ilizarov treatment.



Case Study 32

Antero Lateral Bowing of Left Tibia

- 1,2 & 3 12 years old boy Antero Lateral bowing of left tibia
4. Radiographic view of left tibia before surgery
5. External view of left tibia with Ilizarov fixator in situ
6. Final Radiographic view of the left tibia
7. & 8. Clinical appearance of the patient after the surgery (Front & Back View)





Congenital Shortening of Left Tibia

1. 12 years old girl, 4 cm (LLD) shortening of left tibia, (Front view)
2. 12 yrs old girl, 4 cm LLD shortening of left tibia, (Back view).
3. Galeazzi sign (+) or Alis test (+).
4. 4 cm LLD (left tibia).
5. Radiograph of 4 cm shortening (left side).
6. Before surgery, anxious father (Paediatrician) in OR.
7. Tilting to the left side.
8. Distraction osteogenesis with Ilizarov frame, corticotomy was done in the proximal tibia.
9. Surgeon with smiling patient.
10. Patient with Ilizarov frame.
11. Parents (Father Paediatrician and Mother Skin and VD specialist) are happy with their daughter.
12. Clinical appearance of the patient after correction of LLD (4 cm) (side view).
13. Clinical appearance of the patient after correction of LLD (4 cm) front view.
14. Final follow up after 6 months with the Professor M.M. Bari.



Cont...



Congenital Shortening of Left Tibia (4.8 cm LLD) with Absence of 2 Digits and Short Foot

1. 11 years old girl with congenital shorting (4.8 cm) left tibia, Front view
2. 11 years old girl with congenital shorting (4.8 cm) left tibia, Back view
3. 11 years old girl with congenital shorting (4.8 cm) left tibia, lying position
4. 11 years old girl with congenital shorting (4.8 cm) left tibia, gleazzi sign (+)
5. 11years old girl with congenital shorting (4.8 cm) left tibia, with Ilizarov fixator
6. Radiographic view of left tibia fibula with Ilizarov in situ after 3 months follow up
7. Patient can walk with Ilizarov fixator, almost corrected LLD
8. Clinical appearance of the patient after 7 months follow up



Case Study 35

Normal Height Increase of an Individual Using Ilizarov Technique


1. Former avg. height: 5'0", age 23 years
2. After 7 months of treatment
3. Author is checking the Ilizarov frame
4. After 9 months of treatment
5. After 10 months of treatment
6. Height gained by 4" after 11 months of treatment
7. Previous Height 5'0"
8. New Height: 5'4"



Cont...

Case study 35 radiographic follow up:

1. Normal radiograph of both tibia fibula before lengthening
2. After application of Ilizarov in both the limbs with corticotomy
3. Distraction osteogenesis is seen in both the tibia fibula
4. & 5. Radiographic result after lengthening



Case Study 36


Normal Height Increase

1. & 2. 22 years old girl, normal height 4'8" before surgery,
3. & 4. Placement of introducing Ilizarov wires.
5. After 3 months follow up with Ilizarov apparatus
6. After 6 months follow up with Ilizarov apparatus.
7. Radiographic result after 11 months.
8. & 9. Clinical appearance of the patient, height is increased upto 5'1".



Normal Height Increase

1. 23 Yrs. old girl (Medical Doctor), height before lengthening 4'-8".
2. During treatment with Ilizarov frame in both legs.
3. Follow up of the doctor checking the frame stability.
4. Clinical follow up after 2 months. 2" lengthening was done.
5. Radiographic view of both the tibia fibula with good regeneration after 3 months.
6. Rehabilitation is going on during treatment period stature lengthening.
7. Smiling Dr. with Ilizarov in both the legs after 5 months.
8. Clinical appearance of the Doctor after 6 months.
9. Radiograph of both tibia fibula after 6 months.
10. Final x-ray.
11. Clinical appearance of the Doctor after 11 months; 3" height is increased.



Case Study 38



Post- Polio Residual Deformity - 5.3 cm Shortening

1. 20 years old girl post- polio LLD (5.3 cm) left tibia shortening, flexion contracture of left knee, unstable hip and ankle.
2. Galeazzi sign (+)
3. Ilizarov in the left knee and leg.
4. After 6 months follow up.
5. No LLD and no flexion contracture of the left knee is visible.



Post Polio Residual Deformity - LLD 8 cm

1. 24 years old male, polio residual deformity of left ankle with 8cm LLD.
2. Galeazzi sign (+)
3. 8cm LLD (left) back view.
4. 8cm LLD left front view.
5. Radiographic view of left tibia fibula after 2 months follow up, a good regenerate is seen which is formed by Ilizarov technique.
6. Patient is very happy after 4 months follow up.
7. After full correction 6 months follow up, no LLD (front view).
8. Back view.



Case Study 40

Post Polio Residual Deformity - LLD 5 cm with Unstable Knee, Ankle Facing Forward, Equinocavovarus Deformity.

1. 24 years old male- Post-polio residual deformity; 5 cm LLD, unstable knee, ankle with front heal, equino cavo varus deformity.
2. Back view
3. Walks with hand knee support
4. In silting position
5. In OR
6. During treatment with Ilizarov apparatus. After 6 months follow up. Gradual controlled co-ordinated stretching was done for the correction of the deformity.
7. With thigh, knee, ankle and foot orthosis, patient can walk without any support.
8. Clinical appearance of the patient after 7 months.



Post Polio Residual Deformity, 4 cm LLD With Right Ankle Valgus.

1. 22 years old girl. Post polio residual deformity 4 cm shortening with right ankle valgus.
2. Back view.
3. Front view.
4. Galeazzi sign (+)
5. During treatment with Ilizarov apparatus.
6. Radiograph of right tibia fibula after lengthening.
7. Arthrodesis of right ankle and lengthening of tibia fibula by Ilizarov technique.
8. Clinical appearance of the patient, final outcome after 7 months.
9. Both legs are equal.
10. Clinical appearance of the patient, final outcome after 7 months patient is happy.



Case Study 42


Post Polio Residual Deformity of Right Leg, Ankle and Foot with External Tibial; 2 cm LLD with Equinocavovarus $>65^\circ$ (Severe).

- 19 years old girl. Post polio residual deformity of right leg, ankle and foot with external tibial torsion, 2cm shortening with equinocavovarus deformity $>65^\circ$ degree (severe).
- Back view.
- Lying position.
- Sitting position.
- Sitting position back view.
- Radiograph of right and left tibia 2cm shortening.
- After removal of the Ilizarov apparatus in O.R. Deformity of right leg ankle and foot, external tibial torsion, equino cavo varus deformity corrected.
- Plaster immobilization.
- Radiograph result of right tibia fibula with Ilizarov fixator *in situ*.
- Clinical appearance of the patient after 11 months follow up. (front view).
- Clinical appearance of the patient after 11 months follow up (back view).
- No LLD, right tibia is lengthened 2 cm and torsion is corrected.
- Clinical appearance of the patient, sitting position after 1 year.
- Author with the patient.
- Clinical appearance of the patient after 1 year.



Neurofibromatosis

1. 25 years old male – Neurofibromatosis (Recklen housen’s disease) 5 cm LLD, right side is > left side. 2. Big patch in the lateral chest wall.
2. 5cm LLD.
3. Galeazzi sign (+)
4. 5 cm part of tibia and fibula is removed from the right side.
5. During treatment with Ilizarov apparatus.
6. After removal of Ilizarov fixator (Front view).
7. Back view.
8. & 9. Final follow up after 2 years



Case Study 44


Neurofibromatosis

1. 18 cm LLD neurofibromatosis (von Recklinghausens disease)
2. 9 cm removed from tibia
3. 9 cm removed from femur
4. Visible neuroma
5. After removal of the segments from tibia and femur the whole extremity is fixed with Ilizarov fixator, debulking was also done
6. After treatment.



Fibrous Dysplasia Right Tibia

1. 17 years old boy with procurvatum deformity of right tibia.
2. Procurvatum deformity (Front view).
3. Right tibia with procurvatum deformity (Lateral view).
4. Radiograph of right tibia segmental non union with fibrous dysplasia.
5. Patient with Ilizarov fixator.
6. Radiograph of right tibia fibula with proximal metaphyseal corticotomy and gradual correction of non union site.
7. Close up view of right leg with Ilizarov ring fixator.
8. After 8 months follow up. Radiograph with good regenerate in the corticotomy site and non-union site.
9. Radiographic final result after 9 months follow-up correction deformity and union is achieved.
10. & 11. Clinical appearance of the patient after 9 1/2 months follow up.



Case Study 46

Hypertrophic Nonunion of Tibia, Monkey Mouth Deformity

1. & 2. Radiograph of hypertrophic nonunion of right upper tibia with posteroomedial deformity, monkey mouth deformity.
3. & 4. Clinical appearance of 23 years old boy with posteromedial deformity of right tibia.
5. Radiograph of right tibia fibula with almost corrected deformity by gradual distraction with hinges after 3 months follow up.
6. The right leg is almost straight.
7. Lateral view of right leg.
8. Close up view of right leg with Ilizarov *in situ*.
9. Radiographic result of right tibia fibula after 8 months follow up.
- 10, 11 & 12. After removal of Ilizarov fixator, happy patient is showing "V" sign.
13. & 14. Clinical appearance of the patient after



Cont...



Case Study 47

Post Traumatic Disorganized Knee, Ankle Valgus and 14 cm LLD.

1. 28 years old male- post traumatic right disorganized knee, bad scar in the leg ankle valgus and, 14 cm LLD. Got injury at the age of 7.
2. Patient is in standing position.
3. In the operation theatre.
4. Knee is disorganized. Radiographic view before treatment.
5. During treatment, after 1 month follow up.
6. During treatment, after 2 months follow up.
7. Smiling treatment after 4 months follow up.
8. After 5 months follow up, distraction of the knee is going on.
9. Radiograph of right tibia and femur after 1 month follow up.
10. Radiograph of right tibia and femur after 2 months follow up.
11. Radiograph of tibia and femur after 3 months follow up.
12. A good regenerate is seen in the lower femur after 3 months.
13. A good regenerate is seen after 4 months.
14. After 1 year of follow up (front view).
15. After 1 year of follow up (Back view).
16. After 13 months of follow up.
17. After 14 months of follow up.
18. Clinical appearance of the patient, final follow up after 16 months. (front view).
19. Clinical appearance of the patient, final follow up after 16 months. (Back view).




Cont...



Case Study 48

Mal United Right Upper Tibia, Varus Deformity with 4 cm Shortening with External Rotation Accompanyied by Equinus Deformity >45°

1. External view of the patient, varus deformity with external rotation and 4 cm LLD. (Front view) before surgery.
2. External view of the patient, varus deformity with external rotation and 4 cm LLD. (Back view) before surgery.
3. External view of the patient, varus deformity with external rotation and 4 cm LLD. (sitting position) before surgery.
4. Radiographic view of the same patient, mal union of right upper tibia.
5. Radiographic view of the right tibia during treatment with Ilizarov fixator in situ, 4 telescopic rods with 2 rings are seen. follow up after 4 months.
6. Patient is in Ilizarov fixator in the leg and foot (after 5 months follow up).
7. Radiographic view of the right tibia after 6 months follow up.
8. Clinical appearance of the patient. Final out come after 6 months. (Back view)
9. Clinical appearance of the patient. Final out come after 6 months. (Front view)



Mal Union Right Middle Tibia with Bowing Deformity


1. 29 years old male having right bowing of middle tibia (Frontview) before surgery.
2. 29 years old male having right bowing of middle tibia (Back view) before surgery.
3. In sitting position bowing of middle leg is seen.
4. Radiographic view of mal united right tibia fibula with varus deformity.
5. Radiographic view of corrected varus deformity of right tibia fibula with Ilizarov frame *in situ*.
6. External view of the right leg with Ilizarov frame during treatment, after 4 months follow up.
7. Radiographic result of right tibia fibula after 6 months.
8. Clinical appearance of the patient after 7 months follow up, No Varus deformity is seen.
9. Clinical appearance of the patient after 7 months follow up, patient is in sitting positing. No Varus deformity is seen.



Case Study 50

Genu Valgum Deformity

1. & 2. 26 years old female with left sided genu
2. valgum deformity, 4 cm shortening.
3. Radiographic view of the left knee before surgery.
4. Radiographic view of the left knee with opening wedge osteotomy with visible hinges.
5. & 6. patient with Ilizarov apparatus, almost
6. corrected genu valgum (Front View & Back View).
7. Final corrected radiographic result of the genu valgum of the same patient.
8. & 9. Clinical of the patient, full correction is achieved (Front & Back View).



Genu Valgum Deformity

1. & 2. 9 Years old boy before surgery, left sided genu valgum deformity (Front & Back View).
3. Radiographic view of the Genu Valgum Deformity before surgery.
4. Patient with Ilizarov apparatus after the surgery.
5. Patient is in standing position with Ilizarov apparatus.
6. Clinical appearance of the patient after 4 months follow up.





Chapter XVI

Author with Foreign Dignitaries

Author with Foreign Dignitaries



Prof. O.V. Oganesyanyan is teaching his methodology to Dr. M.M. Bari in CITO, Moscow in May 2003



Prof. G.S. Kulkarni (Ex. IOA President) with Dr. M.M. Bari in Miraj 2003



Ilizarov workshop in BOSCON 2014; Prof. Milind Kulkarni (Miraj), Prof. B.M. Mirazimov (Tashkent), Dr. Ruta Kulkarni (Miraj) & Dr. M.M. Bari, behind Nabia Bari.



Miami Field Hospital, Haiti, February 2010. Author did the Ilizarov surgery of Earth quake victims.



Prof. R.J. Garst (Founding father of NITOR) and Dr. M.M. Bari in BOSCON conference 1999.



BOSCON Feb. 2004, from left to right: Dr. Ruta Kulkarni (Miraj), Dr. M.M. Bari, Prof. B.M. Mirazimov (Tashkent), Prof. Nurul Absar, Prof. K.M. Sirajul Islam.



Prof. M.M. Bari is presenting his to Prof. Dror Paley (USA) on 23.05.2014 in Saint Petersburg.



Prof. R.J. Garst author and Nabia Bari, 1997.



Prof. B.M. Mirazimov (Former Director to Tashkent Scientific Research Institute of Orthopaedics and Traumatology; Ph.D. Supervisor of Prof. M.M. Bari) with Authors family in 2004 at Dhaka.



All Ilizarovians from different countries of the world in Limb deformity course in GOA, April 2005 including Dr. M.M. Bari



Prof. M.M. Bari with his post-doctoral supervisor Academician Prof. V.I. Shevtsov, Kurgan, Russia, 2011



Dr. M.M. Bari is presenting scientific paper on non union of humerus by Ilizarov techniques in Golden Jubilee conference of Indian Orthopaedic Association, December 2005 (Mumbai)

Author as a Faculty in Different Countries of the Globe



6th OASAC Conference, Thimphu, Bhutan 2010



55th IOACON, 2010, Jaipur.



Honored Prof. Ceremony, RISC, RTO Kurgan, Russia, June 2013



Rio de Janeiro, Brazil November 2014



Sign Conference, USA, 2008.



7th International ASAMI Conference, Greece, 2010



APOA Conference, Pattaya, Thailand, 2014



APOA Conference, Delhi, 2012



PAN ARAB Orthopaedic Association Conf. Nov. 2014



Chairman, PAN ARAB Orthopaedic Association Conf., Nov. 2014



8th International ASAMI Conference, GOA, India, 2014



2nd Annual Meeting of Paediatric Orthopaedics, Beijing, China, 2013



SICOT, Rio-de-Jeneiro, 2014



National Ilizarov Workshop- Gorakhpur, UP, India, 2014



34th Annual Conference of North Zone Indian Orthopaedic Association, 2015



APOA Conf. Mumbai, India, 2015



Author with Faculties of ASAMI Russia Conference 23 May, 2014 in Saint Petersburg, Park Inn Hotel.

References:

1. Bari (2013) Severely injured open fractures of long bones: definitive management by Ilizarov method in emergency operation theatre of national institute of traumatology & orthopaedic rehabilitation (NITOR), Dhaka, Bangladesh, Genij Ortopedii 3.
2. Bari MM (2013) A color atlas of limb lengthening surgical reconstruction and deformity correction by Ilizarov Technique 386-395.
3. Bari MM, Islam Shahidul, Shetu NH, Mahfuzer RM (2015) Correction of Leg-Length Discrepancies by Ilizarov Technique, MOJ Orthop Rheumatol 2(1): 00029.
4. Dror Paley (2002) Principles of Deformity correction, Springer-Verlag Berlin Heidelberg.
5. Ilizarov GA (1992) Transosseous Osteosynthesis Theoretical and Clinical Aspects of the Regeneration and Growth of Tissue. Springer-Verlag.
6. Shevtsov VI, Makushin VD, Kufyrev LM (2000) Defects of the Lower Limb Bones. Kurgan, p. 682.
7. Shevtsov VI (2002) Surgical lengthening of the lower limbs / A. V. Popkov. Kurgan, p. 222.
8. Golyakhovsky V, Victor HF (1993) Operative Manual of Ilizarov Technique Ed Mosby.

Glossary

Ambulance: Vehicle designed for the transportation of the sick or injured. They provide immediate emergency care as well as safe and speedy transportation to the hospital. Modern ambulances are equipped with devices for administering blood transfusions, providing pure oxygen, defibrillation and appliances for fracture splinting.

Amputation: This is the surgical removal of a diseased part of and extremity. It is a type of surgery that is done by cutting off or removing infected, dead and harmful part of the body so that the patient's health and function may be preserved or improved. Sometimes it may be caused by trauma or other diseases. These are termed traumatic or auto amputation respectively.

Anesthesia: Absence of physical sensation general or regional.

Anatomical Position: The reference position of the body; standing or lying and facing the observer, with the palms of the hands facing forward.

Anatomical Reduction: The exact adaptation of fracture fragments (hairline adjustment), it will result in complete restoration of the normal anatomy. While overall stability does not depend on precise reduction, precise reduction more reliably results in stability and increased strength of fixation. It is more important in articular fractures than in diaphyseal fractures.

ANGULATION: The orientation of one bone fragment in such a manner that the two parts meet at an angle rather than a straight line. The standard surgical convention is that the angulations are characterized by describing the deviation of the distal part from its anatomical position. For example, in a Colle's fracture, the distal radial fragment is dorsally (or posteriorly) angulated, even though the apex of the deformity points anteriorly; similarly a tibial fracture whose apex angulation points backward should be referred to as angulated anteriorly, as the distal part is indeed angulated anteriorly from its anatomical position.

Ankylosis: Fusion of a joint by bone or a tight fibrous tissue which may occur as a result of a disease process.

Anterior: The front aspect of the body in the anatomical position. If A is in front of B in the anatomical position, then A is said to be anterior to B while B is posterior to A.

Antiseptic: Originally the surgical strategy for avoiding post operative sepsis by applying to the wound bactericidal chemicals, as in the carbolic acid aerosol described and used by Joseph Lister in the late 19th century the era of antiseptic surgery. Now a term used for non biological chemicals which have topical bactericidal properties.

Arthritis: Literally, an inflammatory condition of a synovial joint. It may be septic or aseptic. The former may be blood borne (haematogenous), as in children, or it may follow penetration of the joint by wounding or surgery. Aseptic Arthritis are usually of the rheumatoid type (including Reiter's syndrome, psoriatic arthropathy, etc) or due to degenerative change.

Arthrodesis: Fusion of a joint by bone as a planned outcome of a surgical intervention.

Arthroplasty: Surgical reconstruction or replacement of part or whole joint with or without metal.

Arthrotomy: Cutting into the joint as a form of treatment or diagnosis.

Articular Fracture: Fracture involving the joint as well. It is a serious condition that usually requires open surgical intervention urgently.

Avascular Necrosis (AVN): Bone death due to a cut in its blood supply by disease or injury. This results from injury, infection and metabolic as well as idiopathic diseases. In the absence of sepsis, this is called aseptic necrosis. The dead bone retains its normal strength until the natural process of revascularization by "creeping substitution" starts to remove the dead bone, in preparation for the laying down of new bone. Where the above process fails or there is undue load on the affected areas, segmental collapse takes place. This is commoner in the femoral head and the talus than at other skeletal sites.

Avulsion: Pulling off, e.g. a bone fragment pulled off by a ligament or muscle attachment is an avulsion fracture.

Biomechanics: Is the application of mechanics principles to biologic systems.

Biopsy: The surgical removal of a piece of tissue for histological examination, usually undertaken to establish a diagnosis.

Bone Graft: Bone removed from one skeletal site and placed at another. Bone grafts are used to stimulate bone union and also to restore skeletal continuity where there has been bone loss. Auto graft is a graft of tissue from one site to another within the same individual (homograft) while an allograft is one from another individual of the same species, who is genetically different from the recipient.

Callus: A tissue complex formed at a site of bony repair. Following a fracture it makes a gradual and progressive transition through a series of tissue type haematoma granulation tissue fibrous tissue (or fibro cartilaginous tissue) calcified tissue remodeling into woven bone, gaining in stiffness as it does so, Callus formation is the response of living bone to any irritation (chemical, Infective, mechanical instability, etc. In Internal fixation with absolute stability, where direct (callus free) bone healing is expected the appearance of callus is a sign of unexpected mechanical instability (formerly referred to as irritation) and will alert the surgeon to a failure of the original mechanical objective. Callus is welcome as a repair tissue in all treatment methods where relative fracture stability has been the planned goal.

Cancellous Bone: Is the spongy trabecular bone found mostly at the proximal and distal bone ends in contrast with the dense cortical bone of the shafts. Cancellous bone has a much larger surface area per unit volume and is, therefore, more readily available to its blood supply, as well as to osteoclasts for resorption. Its large surface/volume ratio also offers more surface for invading blood vessels attempting to revascularise dead cancellous bone, and this is an advantage when cancellous bone is used for grafting.

Casts: Casts are materials wrapped over the part of the body to provide immobilization. Mainly used on the extremities or spine following injuries, or in cases of other abnormalities of bone or soft tissues. It can be in the form of plaster of Paris (POP) or fiberglass. Casts when wrapped circumferentially around the extremity, provide more rigid support than splints.

Causalgia: A type of neuralgia or recurrent intractable pain occurring in an extremity as a complication of injury or surgical operation,

Chondrocytes: The active cells of all cartilage, hether particular cartilage, growth cartilage, fibrocartilage etc. they produce the chondral matrix, both its collagen and the muscopolysaccharides of the ground substance.

Coma: A level of consciousness where the patient fails to respond to all level of stimulation.

Compartment Syndrome: Is a condition in which increased pressure within a limited space compromises the circulation and function of the tissues within that space. The major factors in the pathology are:

- Elevated tissue pressure within a closed fascial space
- Reduced tissue perfusion -ischemia
- Leading to cell death – necrosis

Comminution: A condition where a fracture has more than two fragments, such fracture is termed a comminuted fracture.

Consolidation: A stage in bone healing when there is stability and the woven bone (callus) is converted to mature bone.

Corticotomy: It is low energy to osteotomy with the preservation of Periosteum, endosteum and bone marrow.

Cortical Bone: The dense bone forming the tabular element of the shaft, or diaphysss (middle part) of a long bone. The term is also applied to the dense, thin shell covering the cancellous bone of the metaphysis. The term is generally used interchangeably with cortex. Cortical bone which has been completely deprived of its blood supply for any extended period of time dies. It may become revascularized by ingrowth of blood vessels and the newly formed Haversian canals which result from the penetration of osteons.

Damage Control Orthopaedics:

Emergency procedures aimed at rapid reduction and fixation and spanning of periarticular fractures has been termed "damage control orthopaedics".

Debridement: The process of removing dead dying tissue as well as foreign bodies from a wound. Literally, it involves the surgical excision of all avascular, contaminated, infected or other undesirable tissue from

the wound until there is bright red bleeding. Strictly speaking, it refers to the extension of a wound and the opening up of the planes of the injured tissue, usually in the context of open fractures, as described by Ambrose Pare in the 16th century. Today, some centres are using maggots for debridement of infected wounds.

Deformity: Any abnormality of the form or shape of a body part or in other words deviation from normality is called deformity.

Delayed Union: May be defined as a failure of a fracture to consolidate within the normal expected time, which varies according to fracture type and location. Delayed union, like union is a surgical judgment and cannot be allocated time period.

Direct Healing: A type of fracture healing observed with absolutely stable (rigid) internal fixation. It is characterized by:

1. Absence of callus formation specific to the fracture site.
2. Absence of bone surface resorption at the fracture site.
3. Direct bone formation, without any intermediate repair tissue.

Direct fracture healing was formerly called "primary" healing, a term avoided today so as not to imply grading of the quality of fracture healing. Two types of direct healing are distinguished, namely contact healing and gap healing.

Disaster: It is also termed a Major incident. It occurs when the number, severity and type of casualties overwhelm the local medical resources such as personnel, equipment and consumables. It is usually sudden, unexpected and overwhelms the initial response mechanism of the Emergency department. Disaster can be natural or man-made and can occur as, trauma natural disasters, public health disasters or war and civil disorder.

Dislocation: A displacement, usually traumatic of the components of a joint such that no part of one articular surface remains in contact with the other. The term subluxation applies when there is partial contact between the two surfaces.

Displacement: Out of place. A fracture is displaced if the fragments are not perfectly anatomically aligned.

Displacements are:

- Translation (ad latus) - shift.
- Angulation (ad axin).
- Rotation (ad peripherium).
- Length (ad longitudinally).

Distraction Osteogenesis: It is a mechanical induction between bony surfaces that are gradually pulled apart in a controlled number.

RICE for treatment of acute swellings- Rest, apply ice pack, Compress and elevate it.

CSM where you check for the circulation, Sensation and Movement in the effected limb.

Endosteal: The adjective derived from endosteum, which means the interior surface of a bone, e.g. the wall of the medullary cavity.

Epiphysis: The end of long bone which bears the articular component; that fuses with the shaft at the point where it was previously separated by the bone growth plate. It develops from the cartilaginous element between the joint surface and the growth plate.

External Fixation: The technique of skeletal stabilization, which involves the implantation into bone of pins, wires or screws that protrude through the integument and are linked externally by bars, rods or other devices.

Fasciocutaneous:

A term describing tissue flaps which include the skin, the subcutaneous tissues, and the associated deep fascia as a single layer.

Fasciotomy: The surgical division of the investing fascia wall of and osseofascial muscle compartment, usually to release pathologically high intra compartmental pressure in compartment syndrome.

Fibrocartilage: Tissue consisting of elements of cartilage and of fibrous tissue. This may be a normal anatomical entity, such as certain intra-articular structures (menisci, triangular fibrocartilage of the wrist,

the symphysis pubis) or constitute the repair tissue after lesion of the articular cartilage.

Healing: Restoration of the original integrity. The healing process after all bone fracture lasts many years, until internal fracture remodeling subsides. For practical purposes, however, healing is considered to be complete when the bone has regained its normal stiffness and strength.

Health as defined by the WHO is a state of "complete physical mental and social wellbeing and not merely the absence of disease or infirmity" (Physical weakness).

Immobilization: Splinting or resting. The act and methods of stopping a patient or a limb from unwanted movement. Avoid vague use of this word e.g. "immobilize a patient". It is better to specify what is to be immobilized e.g. the ankle joint, the affected limb, the cervical spine or the patient (by admitting the patient in hospital).

Impacted Fracture: A fracture in which the opposing bony surfaces are driven one into the other, resulting often in an inherent fracture stability and usually a degree of angulation.

Implants: Mechanical (artificial) devices inserted into the body to maintain fracture after reduction. Implants are main components of osteosynthesis.

Intramedullary Nail Locked Or Unlocked:

An intramedullary nail provides some degree of stability, mainly as a result of its stiffness. An unlocked nail will allow the fragments to slide together along the nail, the fracture must, therefore be provided with a solid support against shortening. For the treatment of multifragmentary fractures, where there is axial instability (the fear of collapse into a shortened position) the nail can be interlocked above and below the fracture locus to prevent this shortening and also to reduce rotational displacement. This is achieved by locking bolts traversing a locking hole prepared in the nail and passing through the cortex on either side of the nail. If the locking hole is round and matches the size of the locking bolt, then static locking has been achieved. If the locking hole is elongated in the nail's long axis, the possibility of a limited excursion of axial movement is achieved, whilst preserving the rotational control so called dynamic locking.

Ischemia: Pathological absence of blood flow. Inadequate or lack of supply of blood to a part of the body, caused by partial or total blockage of an artery.

Malunion: A fracture that unites in an abnormal anatomical alignment and or rotation. (Consolidation of a fracture in a position of deformity).

Mass Casualty: Generally, a mass casualty situation arises when the number of casualties arriving suddenly at a given time overwhelms the resources of hospital emergency services. These resources include personnel, equipment, infrastructure and consumables. It is also termed "major incident".

Metaphysis: The segment of a long bone located between the end part (epiphysis) and the shaft (diaphysis). It consists mostly of cancellous bone within a thin cortical shell.

Musculoskeletal System: Body system comprising of muscles, bones and all other related tissues. The branch of Medicine that handles the problems of this system and all that makes it to function is called ORTHOPAEDICS. The specific area of Orthopaedic that deals with the study and management of injuries, particularly that of musculoskeletal is called TRAUMATOLOGY.

Neuropraxia: Temporary alteration of nerve functions following ischaemia from a bruising or compression of a peripheral nerve.

Neurotmesis: Damage to the nerve fibre (neuron) and the sheath, a severest form of nerve injury with poor result if not promptly and expertly treated (Nerve transection).

Nonunion: (Author's definition) Arrest of bony fracture repair process with the formation of fibrous or cartilaginous tissue in between the main fragments and fracture has remain un-united for 6-9 months due to mechanical or biological failure judged by clinical and radiological evidence.

Open Fracture: This is a fracture whereby the fragment (s) communicates with the exterior including communication with hollow viscus. This is the opposite of closed fracture. Open fractures are classified into three types namely, types-I, II and III.

Orif: An abbreviation for open reduction and internal fixation. A form of osteosynthesis whereby an implant material is used in the fixation.

Orthopaedics: The word "Orthopaedics" was coined from two Greek words, "Orthos," meaning, "straight" and "paedion", "meaning" "child". This relation of the subject to children arose from the historical fact that orthopaedic practice at that time was based on "straightening-up" of children made "crooked" by various crippling diseases, especially poliomyelitis which was prevalent at that time. This term was coined by Nicholas Andry, a Professor of Medicine at the University of Paris in 1741. Modern Orthopaedics has gone beyond this; it manages all abnormal conditions, including injuries, affecting the musculoskeletal system.

For the Residents Orthopaedics is that branch of surgery that deals with the study and management (Diagnosis, treatment and prevention) of the problems of the musculoskeletal system and its related structures.

Areas Covered In Orthopaedics Include:

- Congenital and developmental problems
- Degenerative (aging) conditions
- Metabolic/endocrine problems of bone & joints
- Tumors and tumor-like conditions
- Injuries and related problems (Traumatology)
- Infection and inflammatory conditions
- Neuromuscular disorders

Orthotics: The branch of medical engineering concerned with the design and fitting of devices such as braces in the treatment of orthopedic disorders. Such devices and appliances are grouped as orthotics.

Osteoarthritis: This is a condition which affects (synovial) joints and is characterized by loss of articular cartilage, reactive subchondral bone sclerosis (sometimes with subchondral cysts), and the formation of peripheral bony outgrowths called osteophytes. The primary lesion is degeneration of the articular cartilage as a consequence of infection, trauma, overuse, congenital skeletal anomaly, or a part of the aging process.

Osteoblast: Cell that form new bone.

Osteoclast: Cell that destroy bone. They are typically responsible for remodeling and are found at the tip of the remodeling osteons. They are also found in all sites where bone is being removed by physiological process.

Osteoclasis: Breaking down (Surgical) a mal united fracture in order to offer the correct treatment. Opposite of osteosynthesis.

Osteomalacia: Insufficient mineralization.

Osteomyelitis: An acute or chronic inflammatory condition affecting bone and its medullary cavity, usually the result of bone infection. This may be a blood borne infection (haematogenous osteomyelitis) usually in children or in the immuno compromised or followed by an open fracture (post traumatic osteomyelitis). The acute form, if diagnosed early and treated vigorously, can heal with no residual effects. If the diagnosis is delayed or treatment neglected, then the infection and the consequent interference with the local vascularity, can result in dead bone (which may separate to form one or more sequestra that remains infected in the long term because the defence mechanisms have no vascular access to it. The treatment of chronic osteomyelitis is surgical and includes wide excision of all dead and infected tissue, the identification of the responsible organism and the delivery, both locally and systemically of appropriate antibacterial agents. In modern world Iliizarov is the best option for chronic osteomyelitis.

Osteon: The name given to the small channels which combine to make up the Haversian system in cortical bone.

Osteopenia: An abnormal reduction in bone mass. This may be generalized, as in some bone disease, or localized, as response to inflammation, infection, disuse, etc. Reduce bone density on Radiograph.

Osteoporosis: A reduction in bone mass in a unit volume (insufficient bone mass). It is a natural aging process but may be pathological. It can result in pathological fracture (Most fractures of the femoral neck in the elderly are due to osteoporosis plus minimal trauma).

Osteosynthesis: This term was coined by Albin Lambotte to describe the fixing of a fractured bone by a surgical intervention using implanted material (surgically "united"). It differs from "internal fixation" in that it also includes external fixation.

Osteotomy: Controlled surgical division of a bone.

Pathological Fracture: A fracture in a diseased bone. It may be the result of the application of a force less than which would be required to produce a fracture in a normal bone.

Periosteal: Relating to or derived from periosteum.

Periosteum: The inelastic membrane bounding the exterior surface of a bone. The periosteum plays an active part in the blood supply to cortical bone, in fracture repair, and in bone remodeling. It is continuous with the perichondrium - the membrane that bounds the periphery of the physis.

Pin Loosening: The pins of external fixator frames serve to stabilize the fragments of a fracture by linking the bone to the frame. Stability depends, among other things, upon the contact between pin and bone (pin loosening occurs when bone surface resorption at the pin bone interface takes place due to excessive cyclical loading of the bone (Note also pin tract infection).

Plaster Of Paris (POP): A white powder, calcium sulfate, that when mixed with water forms a quick hardening paste. It is used in the arts for sculpting and making casts, and in medicine for molding casts around broken limbs.

Pneumothorax: The presence of air in the pleural cavity surrounding the lungs, causing pain and difficulty in breathing. Pneumothorax can occur spontaneously because of accidental rupture or perforation of the pleura, and in the past it was also a deliberate medical procedure in the treatment of tuberculosis.

Polytrauma: Is defined as a clinical state followed by injury to the body leading to profound physio-metabolic changes, involving multiple systems;

In other words:

1. 2 major system and + one major limb injury
2. One major system + 2 major skeletal injuries
3. Unstable fracture pelvis with visceral injury
4. 1 major system injury + 1 open G III skeletal Injury.

Pseudoarthrosis: Means nonunion with neoarthrosis, develops a neoarthrosis with a synovial lining and sometimes with joint fluid in the cleft or cavity.

Prosthesis: The replacement of a missing body part by an artificial substitute is called prosthesis; the term is also used loosely to describe the artificial device; the branch of surgery dealing with prosthesis is prosthetics.

Reduction: The realignment of a displaced fracture.

Refracture: A fracture occurring after the bone has solidly bridges, at a load level otherwise tolerated by normal bone. The resulting fracture line may coincide with the original fracture line, or it may be located remote from the original fracture but within the area of bone that has undergone changes as a result of the fracture and its treatment.

Reflex Sympathetic Dystrophy: See Causalgia and fracture disease.

Remodeling (OF BONE): The process of transformation of external bone shape (external remodeling), or of internal bone structure (internal remodeling), or remodeling of the Haversian system.

Resuscitation: Procedure of reviving an unconscious person. It includes the use of basic and advance trauma life support. Such procedures during the first 6 hours (Golden hour) following injury is crucial.

Rigid Fixation: A method of internal fixation where micromotion of the fracture ends is virtually eliminated (concept of Relative stability).

Segmental Fracture: This is a type of fracture where the shaft of a bone is broken at two levels, leaving a separate shaft segment between the two fracture sites.

Sequestrum: A piece of dead bone lying alongside, but separated from, the osseous bone from which it came. It is formed when a section

of bone is deprived of its blood supply. A sequestrum may be aseptic (sterile), as for example, beneath a plate when there has been massive periosteal stripping and then a plate with a high contact "footprint" applied, killing the underlying bone. This is especially seen if a plate has been applied to the cortex at the same time as a reamed intramedullary nail has been inserted. It is a diagnostic feature in chronic osteomyelitis.

Shock: A state of reduced tissue perfusion, usually due to a fall of intravascular pressure secondary to hypovolaemia, overwhelming sepsis (gram negative shock) or allergic anaphylaxis.

Shock-Spinal: A state of temporary failure of the spinal cord functions in the immediate post-injury period. It is characterized by flaccid paralysis, absent reflexes and loss of sensation below the level of the lesion. The condition usually lasts between 24-48 hours, the return of the anal wink and bulbocavernosus reflexes are the hallmarks of recovery.

Spica: A bandage or cast applied to a limb which also incorporates a joint and trunk or applied in an overlapping figure eight pattern to immobilize it, e.g. hip or shoulder spica.

Splint: Support / device to immobilize broken bones rigid material used to keep a broken bone or other injured part from moving.

Splinting: Reducing the mobility at a fracture site by applying a firm object (splint) to the main bone fragments. The splint may be external (plaster, external fixators) or internal (plate, intramedullary nail).

Sprain: This is any painful wrenching (twisting or pulling) movement of a joint ligament. It refers to a less severe injury than capsular or ligamentous tear (ref. strain). Sprain affect joint while strain affects muscle.

Stabilization: Means of making the injured person clinically better or an injured limb stop moving abnormally.

Traction: A device that applies a pulling force through the part of the body aimed at reducing and splinting a fracture or dislocation (skin, skeletal); In other words, directional pull on a trunk or an extremity is called traction.

Trauma: Injury or processes that are injurious to health. It may be physical, emotional, psycho social or economical.

Traumatology: That branch of medical science (Surgery) that deals with treatment and prevention of injuries.

Triage: The process of prioritizing sick or injured people for treatment according to the seriousness of the condition or injury.

Union: Union means "as one" as in marital union. When a fracture is fixed so that the bone functions as a single unit, this does not mean that the bone has healed. Bone healing is a gradual process which continues until the bone is restored to its final state by remodeling which may take years. Fracture union means that a healing process has reached the point when the surgeon guesses that it can withstand normal functional loads for that patient. It is usually based on clinical and imaging information. That is why it is wise not to fix the "time of fracture union" but only estimate it.

Valgus: Deviation away from the midline in the anatomical position. Thus, genu valgum is a deformity at the knee where the lower leg is angled away from the midline (knock knee). By convention, any deformity or deviation is described in terms of the movement of the distal part.

Varus: Deviation towards the midline in the anatomical position. Thus, genu varum is a deformity at the knee where the lower leg is angled towards the midline (bow leg). By convention, any deformity, or deviation, is described in terms of the movement of the distal part.

Special guidelines for (D. Ortho, MS Ortho)

Residents:

1. Prepare your mind to succeed.
2. Acquire the necessary book and tools and use them properly.
3. Learn basic skills and practice them even on your class mates daily.
4. Do not under-rate any posting because all of them are essential for your genuine qualification.
5. Familiarize yourself with each topic (disease/condition) and form notes around them. This is because you will come across them at different levels of your training. For example topics like osteomyelitis or sickle cell anaemia will confront you in all the specialities of Medicine and Surgery.
6. Maintain dairy and personal notes and update them as you pass through the different departments.
7. Do not try to be too familiar with lecturers than your books and patients.
8. Do not allow your parents/guardians to influence any stage of your training because they will not be there when you face the bigger problem associated with patients' care.
9. The profession of orthopaedic and trauma is a lifelong challenge, parents/teachers and counselors should help students to develop self-determination and boldness based on firm knowledge of the subject (even at the admission level). This is important if we want to produce Doctors and not gamblers.

Resit examination should not scare you; rather it should challenge you to work harder. There is no place for part-time study in Medicine and you may not get all the time for cultism and unnecessary social activities. During the study of the basic sciences of Anatomy, Physiology and Biochemistry, students should note the clinical relevance of every topic.

Respect your teachers, parents and love your patients. Peruse your books to encourage those teachers who may want to spend their private time in teaching you.

Fear Almighty Allah

"MedCrave Group is an open access publishing group that caters to those who wish to communicate their scientific ideas and research to the rest of the world. MedCrave provides users with unlimited scientific research information to bolster their research and general knowledge base. On MedCrave, you will find scientific research information on just about every topic you can imagine. Not only do users have the ability to access and view every research paper in MedCrave, but users can also download and share research papers. The primary goal of the MedCrave Group is to make scientific content accessible to the entire global population, and the MedCrave Group has taken many steps to make this goal more achievable."




Why Publishing with MedCrave?

- * Global scientific network with 70+Peer Reviewed Open Access Journals and 4000+ Editorial Board
- * More than 3000 Open Access articles published
- * New initiative in E Books with ISBN & Barcode
- * Global standards (Cover lamination) in Books & Articles Printing
- * More than 5000 Unique online viewers a day

Cordial Invitation for Manuscripts

We would be please to welcome you as one of our authors by submitting valuable manuscripts towards the journal at <http://medcraveonline.com/submitManuscript>

Please visit the journal at <http://medcraveonline.com/>



MedCrave

Step into the World of Research

MedCrave Group

820 W, Danforth Rd.,

Edmond, OK 73003

Tel: +1-918-208-0288

Fax: +1-918-512-4618

Email: info@medcraveonline.com

ISBN 978-0-9967956-1-6

